



10/772,768

01/31/05

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b> <b>C12N 5/06, A61K 35/14,</b> <b>A61K 39/00, C12N 5/08,</b> <b>G01N 33/50</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/00587</b> <b>(43) International Publication Date:</b> 06 January 2000 (06.01.2000)
---	-----------	--

**(21) International Application Number:** PCT/DK99/00363  
**(22) International Filing Date:** 25 June 1999 (25.06.1999)

**(30) Priority Data:**  
60/091,684 02 July 1998 (02.07.1998) US  
PA 1998 00848 26 June 1998 (26.06.1998) DK  
PA 1998 00895 01 July 1998 (01.07.1998) DK

**(60) Parent Application or Grant**  
KALTOFT, Keld [/]; (). AGNHOLT, Jorgen [/];  
(). KALTOFT, Keld [/]; (). AGNHOLT, Jorgen [/];  
(). HOFMAN-BANG & BOUTARD, LEHMANN & REE  
A/S; ().

Published

**(54) Title: METHODS OF EXPANDING AND SELECTING DISEASE ASSOCIATED T-CELLS**  
**(54) Titre: METHODE DE DEVELOPPEMENT ET DE SELECTION DES LYMPHOCYTES T ASSOCIES A UNE PATHOLOGIE**

**(57) Abstract**

Methods of expanding and selecting disease associated T-cells, continuous T-cell lines as well as T-cell lines obtainable by these methods are disclosed. Furthermore, pharmaceutical compositions and vaccines comprising activated disease associated T-cell are disclosed. The uses of the T-cell and T-cell lines are numerous and include methods of diagnosis, methods for the treatment, alleviation or prevention of diseases associated with activation of T-cells, methods of testing the effect of medicaments against T-cell associated diseases, methods of detecting T-cell growth factors, methods of monitoring the response to treatment, alleviation or prevention of diseases associated with activation of T-cells, and methods of identifying disease associated antigens.

**(57) Abrégé**

L'invention porte sur des méthodes de développement et de sélection des lymphocytes T associés à une pathologie de lignées continues de lymphocytes T, ainsi que sur les lignées de lymphocytes T pouvant être obtenues par ces procédés. L'invention porte également sur des compositions pharmaceutiques et des vaccins comprenant des lymphocytes T activés, associées à une pathologie. Les utilisations des lymphocytes T et des lignées de lymphocytes T sont nombreuses et comprennent des méthodes de diagnostic, de traitement, de soulagement ou de prévention de pathologies associées à l'activation des lymphocytes T, des méthodes visant à tester les effets des médicaments permettant de lutter contre des pathologies associées aux lymphocytes T, des méthodes de détection des facteurs de croissance des lymphocytes T, des méthodes visant à surveiller les réactions au traitement, le soulagement ou la prévention des pathologies associées à l'activation des lymphocytes T, et des méthodes d'identification d'antigènes associés à des pathologies.

Best Available Copy

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 5/06, 5/08, A61K 39/00, 35/14,</b> <b>G01N 33/50</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/00587</b> <b>(43) International Publication Date:</b> 6 January 2000 (06.01.00)
<b>(21) International Application Number:</b> PCT/DK99/00363 <b>(22) International Filing Date:</b> 25 June 1999 (25.06.99) <b>(30) Priority Data:</b> PA 1998 00848 26 June 1998 (26.06.98) DK PA 1998 00895 1 July 1998 (01.07.98) DK 60/091,684 2 July 1998 (02.07.98) US <b>(71)(72) Applicants and Inventors:</b> KALTOFT, Keld [DK/DK]; Voldby Hovvej 10, Voldby, DK-8450 Hammel (DK). AGN- HOLT, Jørgen [DK/DK]; Temevej 12, DK-8240 Risskov (DK). <b>(74) Agent:</b> HOFMAN-BANG & BOUTARD, LEHMANN & REE A/S; Hans Bekkevolds Allé 7, DK-2900 Hellerup (DK).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>	
<b>(54) Title:</b> METHODS OF EXPANDING AND SELECTING DISEASE ASSOCIATED T-CELLS  <b>(57) Abstract</b>  Methods of expanding and selecting disease associated T-cells, continuous T-cell lines as well as T-cell lines obtainable by these methods are disclosed. Furthermore, pharmaceutical compositions and vaccines comprising activated disease associated T-cell are disclosed. The uses of the T-cell and T-cell lines are numerous and include methods of diagnosis, methods for the treatment, alleviation or prevention of diseases associated with activation of T-cells, methods of testing the effect of medicaments against T-cell associated diseases, methods of detecting T-cell growth factors, methods of monitoring the response to treatment, alleviation or prevention of diseases associated with activation of T-cells, and methods of identifying disease associated antigens.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**Description**

5

10

15

20

25

30

35

40

45

50

55

5

METHODS OF EXPANDING AND SELECTING DISEASE ASSOCIATED T-CELLS

10

15

20

25

30

35

40

45

50

55

5 The present invention relates to methods of expanding and  
selecting disease associated T-cells, continuous T-cell  
lines as well as T-cell lines obtainable by the methods.  
The invention also relates to pharmaceutical compositions  
comprising activated disease associated T-cell. In a  
10 further aspect, the invention relates to vaccines  
comprising such activated disease associated inflammatory  
T-cells. The invention further relates to pharmaceutical  
compositions for use in adjuvant treatment comprising  
disease associated regulatory or cytotoxic T-cells.  
25 Furthermore, the present invention concerns the use of T-  
cell lines for preparing medicaments for treating T-cell  
associated diseases as well as for use in a broad range  
of methods, i.a. methods of diagnosis, methods for the  
treatment, alleviation or prevention of diseases  
30 associated with activation of T-cells, methods of testing  
the effect of medicaments against T-cell associated  
diseases, methods of detecting T-cell growth factors,  
methods of monitoring the response to treatment,  
alleviation or prevention of diseases associated with  
35 activation of T-cells, and methods of identifying disease  
associated antigens. The present invention also concerns  
a model system for testing the effect of a medicament  
against a T-cell associated disease.

30 BACKGROUND OF THE INVENTION

All normal somatic cells are believed to have a finite in  
vitro life-span commonly known as the Hayflick limit.  
This dogma is a cornerstone in cell biology. According to  
50 this, only a certain number of cell population doublings  
(PD) is possible. Following approximately 23 PD, T-cells

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

go into replicative senescence, and the cells cease to divide. This implies that one T-cell can on average expand only to  $2^{23}$  cells corresponding to approximately  $10^7$  T-lymphocytes. Most often  $10^7$  T-lymphocytes, that is about 10 mg, are not enough "material" for use as T-cell vaccine in treatment of patients with T-cell-related auto-immune/chronic inflammatory diseases or for the use as T-cell adjuvant therapy in patients with inflammatory/auto-immune or malignant diseases. By way of example, in cancer 10 mg of clonal cytotoxic T-cells is far to little to combat tumour masses in the order of kilograms.

10  
15  
20  
25  
30  
35  
40  
45  
50  
55

In the prior art, there is several examples of attempts to overcome this problem. However, none has come up with the solution presented in the present invention. Several publications relate to activated T-cells wherein antigen specific T-cells are produced ex vivo after stimulation in vitro with a known antigen. The T-cells are commonly produced from peripheral blood T-cells by procedures, in which an antigen is used to stimulate T-cells. The antigen specific T-cell clones are obtained by using conventional immunological selection techniques. Only a few successful attempts to produce disease-associated T-cells in sufficient amount have been reported.

10  
15  
20  
25  
30  
35  
40  
45  
50  
55

WO 88/07077 (Liu) (ref. 1) discloses a method of expanding helper T-cells ( $T_h$ -cells) recognising viral antigens, wherein  $T_h$  cells are made to proliferate from a sample of mononuclear cells including the  $T_h$ -cells and antigen presenting cells (APCs) by the addition of specific viral antigen. The proliferating  $T_h$ -cells may be expanded in the presence of APCs and specific antigen. Optionally IL-2 may be added in order to stimulate the expansion.

5  
10  
15  
20  
25  
WO 94/02156 (Engelman) (ref. 2) discloses a method of activating T-cell isolated from peripheral blood, wherein specific antigen is used to pulse dendritic cells and thereafter mixed with the isolated T-cells. The mixture is expanded in the presence of IL-2 and /or IL-4, however, in very low concentrations (about 2 IU/ml).

30  
35  
40  
45  
50  
55  
WO 97/05239 (Gruenberg) (ref. 3) discloses a method of expanding T-cells isolated from the peripheral blood, wherein the expansion is performed without IL-2 due to its alleged toxic effect in humans.

60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995  
1000  
1005  
1010  
1015  
1020  
1025  
1030  
1035  
1040  
1045  
1050  
1055  
1060  
1065  
1070  
1075  
1080  
1085  
1090  
1095  
1100  
1105  
1110  
1115  
1120  
1125  
1130  
1135  
1140  
1145  
1150  
1155  
1160  
1165  
1170  
1175  
1180  
1185  
1190  
1195  
1200  
1205  
1210  
1215  
1220  
1225  
1230  
1235  
1240  
1245  
1250  
1255  
1260  
1265  
1270  
1275  
1280  
1285  
1290  
1295  
1300  
1305  
1310  
1315  
1320  
1325  
1330  
1335  
1340  
1345  
1350  
1355  
1360  
1365  
1370  
1375  
1380  
1385  
1390  
1395  
1400  
1405  
1410  
1415  
1420  
1425  
1430  
1435  
1440  
1445  
1450  
1455  
1460  
1465  
1470  
1475  
1480  
1485  
1490  
1495  
1500  
1505  
1510  
1515  
1520  
1525  
1530  
1535  
1540  
1545  
1550  
1555  
1560  
1565  
1570  
1575  
1580  
1585  
1590  
1595  
1600  
1605  
1610  
1615  
1620  
1625  
1630  
1635  
1640  
1645  
1650  
1655  
1660  
1665  
1670  
1675  
1680  
1685  
1690  
1695  
1700  
1705  
1710  
1715  
1720  
1725  
1730  
1735  
1740  
1745  
1750  
1755  
1760  
1765  
1770  
1775  
1780  
1785  
1790  
1795  
1800  
1805  
1810  
1815  
1820  
1825  
1830  
1835  
1840  
1845  
1850  
1855  
1860  
1865  
1870  
1875  
1880  
1885  
1890  
1895  
1900  
1905  
1910  
1915  
1920  
1925  
1930  
1935  
1940  
1945  
1950  
1955  
1960  
1965  
1970  
1975  
1980  
1985  
1990  
1995  
2000  
2005  
2010  
2015  
2020  
2025  
2030  
2035  
2040  
2045  
2050  
2055  
2060  
2065  
2070  
2075  
2080  
2085  
2090  
2095  
2100  
2105  
2110  
2115  
2120  
2125  
2130  
2135  
2140  
2145  
2150  
2155  
2160  
2165  
2170  
2175  
2180  
2185  
2190  
2195  
2200  
2205  
2210  
2215  
2220  
2225  
2230  
2235  
2240  
2245  
2250  
2255  
2260  
2265  
2270  
2275  
2280  
2285  
2290  
2295  
2300  
2305  
2310  
2315  
2320  
2325  
2330  
2335  
2340  
2345  
2350  
2355  
2360  
2365  
2370  
2375  
2380  
2385  
2390  
2395  
2400  
2405  
2410  
2415  
2420  
2425  
2430  
2435  
2440  
2445  
2450  
2455  
2460  
2465  
2470  
2475  
2480  
2485  
2490  
2495  
2500  
2505  
2510  
2515  
2520  
2525  
2530  
2535  
2540  
2545  
2550  
2555  
2560  
2565  
2570  
2575  
2580  
2585  
2590  
2595  
2600  
2605  
2610  
2615  
2620  
2625  
2630  
2635  
2640  
2645  
2650  
2655  
2660  
2665  
2670  
2675  
2680  
2685  
2690  
2695  
2700  
2705  
2710  
2715  
2720  
2725  
2730  
2735  
2740  
2745  
2750  
2755  
2760  
2765  
2770  
2775  
2780  
2785  
2790  
2795  
2800  
2805  
2810  
2815  
2820  
2825  
2830  
2835  
2840  
2845  
2850  
2855  
2860  
2865  
2870  
2875  
2880  
2885  
2890  
2895  
2900  
2905  
2910  
2915  
2920  
2925  
2930  
2935  
2940  
2945  
2950  
2955  
2960  
2965  
2970  
2975  
2980  
2985  
2990  
2995  
3000  
3005  
3010  
3015  
3020  
3025  
3030  
3035  
3040  
3045  
3050  
3055  
3060  
3065  
3070  
3075  
3080  
3085  
3090  
3095  
3100  
3105  
3110  
3115  
3120  
3125  
3130  
3135  
3140  
3145  
3150  
3155  
3160  
3165  
3170  
3175  
3180  
3185  
3190  
3195  
3200  
3205  
3210  
3215  
3220  
3225  
3230  
3235  
3240  
3245  
3250  
3255  
3260  
3265  
3270  
3275  
3280  
3285  
3290  
3295  
3300  
3305  
3310  
3315  
3320  
3325  
3330  
3335  
3340  
3345  
3350  
3355  
3360  
3365  
3370  
3375  
3380  
3385  
3390  
3395  
3400  
3405  
3410  
3415  
3420  
3425  
3430  
3435  
3440  
3445  
3450  
3455  
3460  
3465  
3470  
3475  
3480  
3485  
3490  
3495  
3500  
3505  
3510  
3515  
3520  
3525  
3530  
3535  
3540  
3545  
3550  
3555  
3560  
3565  
3570  
3575  
3580  
3585  
3590  
3595  
3600  
3605  
3610  
3615  
3620  
3625  
3630  
3635  
3640  
3645  
3650  
3655  
3660  
3665  
3670  
3675  
3680  
3685  
3690  
3695  
3700  
3705  
3710  
3715  
3720  
3725  
3730  
3735  
3740  
3745  
3750  
3755  
3760  
3765  
3770  
3775  
3780  
3785  
3790  
3795  
3800  
3805  
3810  
3815  
3820  
3825  
3830  
3835  
3840  
3845  
3850  
3855  
3860  
3865  
3870  
3875  
3880  
3885  
3890  
3895  
3900  
3905  
3910  
3915  
3920  
3925  
3930  
3935  
3940  
3945  
3950  
3955  
3960  
3965  
3970  
3975  
3980  
3985  
3990  
3995  
4000  
4005  
4010  
4015  
4020  
4025  
4030  
4035  
4040  
4045  
4050  
4055  
4060  
4065  
4070  
4075  
4080  
4085  
4090  
4095  
4100  
4105  
4110  
4115  
4120  
4125  
4130  
4135  
4140  
4145  
4150  
4155  
4160  
4165  
4170  
4175  
4180  
4185  
4190  
4195  
4200  
4205  
4210  
4215  
4220  
4225  
4230  
4235  
4240  
4245  
4250  
4255  
4260  
4265  
4270  
4275  
4280  
4285  
4290  
4295  
4300  
4305  
4310  
4315  
4320  
4325  
4330  
4335  
4340  
4345  
4350  
4355  
4360  
4365  
4370  
4375  
4380  
4385  
4390  
4395  
4400  
4405  
4410  
4415  
4420  
4425  
4430  
4435  
4440  
4445  
4450  
4455  
4460  
4465  
4470  
4475  
4480  
4485  
4490  
4495  
4500  
4505  
4510  
4515  
4520  
4525  
4530  
4535  
4540  
4545  
4550  
4555  
4560  
4565  
4570  
4575  
4580  
4585  
4590  
4595  
4600  
4605  
4610  
4615  
4620  
4625  
4630  
4635  
4640  
4645  
4650  
4655  
4660  
4665  
4670  
4675  
4680  
4685  
4690  
4695  
4700  
4705  
4710  
4715  
4720  
4725  
4730  
4735  
4740  
4745  
4750  
4755  
4760  
4765  
4770  
4775  
4780  
4785  
4790  
4795  
4800  
4805  
4810  
4815  
4820  
4825  
4830  
4835  
4840  
4845  
4850  
4855  
4860  
4865  
4870  
4875  
4880  
4885  
4890  
4895  
4900  
4905  
4910  
4915  
4920  
4925  
4930  
4935  
4940  
4945  
4950  
4955  
4960  
4965  
4970  
4975  
4980  
4985  
4990  
4995  
5000  
5005  
5010  
5015  
5020  
5025  
5030  
5035  
5040  
5045  
5050  
5055  
5060  
5065  
5070  
5075  
5080  
5085  
5090  
5095  
5100  
5105  
5110  
5115  
5120  
5125  
5130  
5135  
5140  
5145  
5150  
5155  
5160  
5165  
5170  
5175  
5180  
5185  
5190  
5195  
5200  
5205  
5210  
5215  
5220  
5225  
5230  
5235  
5240  
5245  
5250  
5255  
5260  
5265  
5270  
5275  
5280  
5285  
5290  
5295  
5300  
5305  
5310  
5315  
5320  
5325  
5330  
5335  
5340  
5345  
5350  
5355  
5360  
5365  
5370  
5375  
5380  
5385  
5390  
5395  
5400  
5405  
5410  
5415  
5420  
5425  
5430  
5435  
5440  
5445  
5450  
5455  
5460  
5465  
5470  
5475  
5480  
5485  
5490  
5495  
5500  
5505  
5510  
5515  
5520  
5525  
5530  
5535  
5540  
5545  
5550  
5555  
5560  
5565  
5570  
5575  
5580  
5585  
5590  
5595  
5600  
5605  
5610  
5615  
5620  
5625  
5630  
5635  
5640  
5645  
5650  
5655  
5660  
5665  
5670  
5675  
5680  
5685  
5690  
5695  
5700  
5705  
5710  
5715  
5720  
5725  
5730  
5735  
5740  
5745  
5750  
5755  
5760  
5765  
5770  
5775  
5780  
5785  
5790  
5795  
5800  
5805  
5810  
5815  
5820  
5825  
5830  
5835  
5840  
5845  
5850  
5855  
5860  
5865  
5870  
5875  
5880  
5885  
5890  
5895  
5900  
5905  
5910  
5915  
5920  
5925  
5930  
5935  
5940  
5945  
5950  
5955  
5960  
5965  
5970  
5975  
5980  
5985  
5990  
5995  
6000  
6005  
6010  
6015  
6020  
6025  
6030  
6035  
6040  
6045  
6050  
6055  
6060  
6065  
6070  
6075  
6080  
6085  
6090  
6095  
6100  
6105  
6110  
6115  
6120  
6125  
6130  
6135  
6140  
6145  
6150  
6155  
6160  
6165  
6170  
6175  
6180  
6185  
6190  
6195  
6200  
6205  
6210  
6215  
6220  
6225  
6230  
6235  
6240  
6245  
6250  
6255  
6260  
6265  
6270  
6275  
6280  
6285  
6290  
6295  
6300  
6305  
6310  
6315  
6320  
6325  
6330  
6335  
6340  
6345  
6350  
6355  
6360  
6365  
6370  
6375  
6380  
6385  
6390  
6395  
6400  
6405  
6410  
6415  
6420  
6425  
6430  
6435  
6440  
6445  
6450  
6455  
6460  
6465  
6470  
6475  
6480  
6485  
6490  
6495  
6500  
6505  
6510  
6515  
6520  
6525  
6530  
6535  
6540  
6545  
6550  
6555  
6560  
6565  
6570  
6575  
6580  
6585  
6590  
6595  
6600  
6605  
6610  
6615  
6620  
6625  
6630  
6635  
6640  
6645  
6650  
6655  
6660  
6665  
6670  
6675  
6680  
6685  
6690  
6695  
6700  
6705  
6710  
6715  
6720  
6725  
6730  
6735  
6740  
6745  
6750  
6755  
6760  
6765  
6770  
6775  
6780  
6785  
6790  
6795  
6800  
6805  
6810  
6815  
6820  
6825  
6830  
6835  
6840  
6845  
6850  
6855  
6860  
6865  
6870  
6875  
6880  
6885  
6890  
6895  
6900  
6905  
6910  
6915  
6920  
6925  
6930  
6935  
6940  
6945  
6950  
6955  
6960  
6965  
6970  
6975  
6980  
6985  
6990  
6995  
7000  
7005  
7010  
7015  
7020  
7025  
7030  
7035  
7040  
7045  
7050  
7055  
7060  
7065  
7070  
7075  
7080  
7085  
7090  
7095  
7100  
7105  
7110  
7115  
7120  
7125  
7130  
7135  
7140  
7145  
7150  
7155  
7160  
7165  
7170  
7175  
7180  
7185  
7190  
7195  
7200  
7205  
7210  
7215  
7220  
7225  
7230  
7235  
7240  
7245  
7250  
7255  
7260  
7265  
7270  
7275  
7280  
7285  
7290  
7295  
7300  
7305  
7310  
7315  
7320  
7325  
7330  
7335  
7340  
7345  
7350  
7355  
7360  
7365  
7370  
7375  
7380  
7385  
7390  
7395  
7400  
7405  
7410  
7415  
7420  
7425  
7430  
7435  
7440  
7445  
7450  
7455  
7460  
7465  
7470  
7475  
7480  
7485  
7490  
7495  
7500  
7505  
7510  
7515  
7520  
7525  
7530  
7535  
7540  
7545  
7550  
7555  
7560  
7565  
7570  
7575  
7580  
7585  
7590  
7595  
7600  
7605  
7610  
7615  
7620  
7625  
7630  
7635  
7640  
7645  
7650  
7655  
7660  
7665  
7670  
7675  
7680  
7685  
7690  
7695  
7700  
7705  
7710  
7715  
7720  
7725  
7730  
7735  
7740  
7745  
7750  
7755  
7760  
7765  
7770  
7775  
7780  
7785  
7790  
7795  
7800  
7805  
7810  
7815  
7820  
7825  
7830  
7835  
7840  
7845  
7850  
7855  
7860  
7865  
7870  
7875  
7880  
7885  
7890  
7895  
7900  
7905  
7910  
7915  
7920  
7925  
7930  
7935  
7940  
7945  
7950  
7955  
7960  
7965  
7970  
7975  
7980  
7985  
7990  
7995  
8000  
8005  
8010  
8015  
8020  
8025  
8030  
8035  
8040  
8045  
8050  
8055  
8060  
8065  
8070  
8075  
8080  
8085  
8090  
8095  
8100  
8105  
8110  
8115  
8120  
8125  
8130  
8135  
8140  
8145  
8150  
8155  
8160  
8165  
8170  
8175  
8180  
8185  
8190  
8195  
8200  
8205  
8210  
8215  
8220  
8225  
8230  
8235  
8240  
8245  
8250  
8255  
8260  
8265  
8270  
8275  
8280  
8285  
8290  
8295  
8300  
8305  
8310  
8315  
8320  
8325  
8330  
8335  
8340  
8345  
8350  
8355  
8360  
8365  
8370  
8375  
8380  
8385  
8390  
8395  
8400  
8405  
8410  
8415  
8420  
8425  
8430  
8435  
8440  
8445  
8450  
8455  
8460  
8465  
8470  
8475  
8480  
8485  
8490  
8495  
8500  
8505  
8510  
8515  
8520  
8525  
8530  
8535  
8540  
8545  
8550  
8555  
8560  
8565  
8570  
8575  
8580  
8585  
8590  
8595  
8600  
8605  
8610  
8615  
8620  
8625  
8630  
8635  
8640  
8645  
8650  
8655  
8660  
8665  
8670  
8675  
8680  
8685  
8690  
8695  
8700  
8705  
8710  
8715  
8720  
8725  
8730  
8735  
8740  
8745  
8750  
8755  
8760  
8765  
8770  
8775  
8780  
8785  
8790  
8795  
8800  
8805  
8810  
8815  
8820  
8825  
8830  
8835  
8840  
8845  
8850  
8855  
8860  
8865  
8870  
8875  
8880  
8885  
8890  
8895  
8900  
8905  
8910  
8915  
8920  
8925  
8930  
8935  
8940  
8945  
8950  
8955  
8960  
8965  
8970  
8975  
8980  
8985  
8990  
8995  
9000  
9005  
9010  
9015  
9020  
9025  
9030  
9035  
9040  
9045  
9050  
9055  
9060  
9065  
9070  
9075  
9080  
9085  
9090  
9095  
9100  
9105  
9110  
9115  
9120  
9125  
9130  
9135  
9140  
9145  
9150  
9155  
9160  
9165  
9170  
9175  
9180  
9185  
9190  
9195  
9200  
9205  
9210  
9215  
9220  
9225  
9230  
9235  
9240  
9245  
9250  
9255  
9260  
9265  
9270  
9275  
9280  
9285  
9290  
9295  
9300  
9305  
9310  
9315  
9320  
9325  
9330  
9335  
9340  
9345  
9350  
9355  
9360  
9365  
9370  
9375  
9380  
9385  
9390  
9395  
9400  
9405  
9410  
9415  
9420  
9425  
9430  
9435  
9440  
9445  
9450  
9455  
9460  
9465  
9470  
9475  
9480  
9485  
9490  
9495  
9500  
9505  
9510  
9515  
9520  
9525  
9530  
9535  
9540  
9545  
9550  
9555  
9560  
9565  
9570  
9575  
9580  
9585  
9590  
9595  
9600  
9605  
9610  
9615  
9620  
9625  
9630  
9635  
9640  
9645  
9650  
9655  
9660  
9665  
9670  
9675  
9680  
9685  
9690  
9695  
9700  
9705  
9710  
9715  
9720  
9725  
9730  
9735  
9740  
9745  
9750  
9755  
9760  
9765  
9770  
9775  
9780  
9785  
9790  
9795  
9800  
9805  
9810  
9815  
9820  
9825  
9830  
9835  
9840  
9845  
9850  
9855  
9860  
9865  
9870  
9875  
9880  
9885  
9890  
9895  
9900  
9905  
9910  
9915  
9920  
9925  
9930  
9935  
9940  
9945  
9950  
9955  
9960  
9965  
9970  
9975  
9980  
9985  
9990  
9995  
10000  
10005  
10010  
10015  
10020  
10025  
10030  
10035  
10040  
10045  
10050  
10055  
10060  
10065  
10070  
10075  
10080  
10085  
10090  
10095  
10100  
10105  
10110  
10115  
10120  
10125  
10130  
10135  
10140  
10145  
10150  
10155  
10160  
10165  
10170  
10175  
10180  
1018

5 associated/specific T cells participating in a regulatory  
10 network. The activity of inflammatory T-cells (IFN $\gamma$  and  
TNF $\alpha$  producing) is regulated by IL-10 producing  
regulatory T cells (In a type 1 inflammatory process, the  
5 inverse in type 2 inflammatory processes), cf. Fig. 1.

15 In human studies, it has been very difficult to obtain  
the relevant auto-reactive T-cells and propagate these  
cells into sufficient amounts to produce T-cell vaccines,  
10 although T-cell vaccination studies in disseminated  
20 sclerosis has been promising.

25 Surprisingly, it has now been recognised that continuous  
T-cell lines are obtainable by a method of expanding and  
15 selecting disease associated T-cells. The principle of  
the present invention is based on in vivo antigen  
stimulation, this in vivo stimulation leading to the  
30 presence of a certain population of activated T-cells,  
and this T-cell population can be expanded and selected  
20 under certain conditions. T-cells associated with the  
manifestations of a disease are activated in vivo, and,  
35 may therefore often be expanded in vitro without further  
supplement of a disease associated antigen. Furthermore,  
the T-cells are activated in vivo in such a manner that  
25 they are able to grow in vitro under special conditions.  
40 No cloning step is necessary. The activated T-cells are  
ready to expand and may therefore outgrow non-activated  
T-cells. The pool of activated T-cells in a biopsy can  
45 contain T-cells with different specificities and  
30 functions as well as being of different phenotypes.  
Selection of a T-cell line with a desired phenotype,  
specificity and function may be controlled by the  
50 conditions of the growth media, and by immunoselection  
methods.

55  
35 SUMMARY OF THE INVENTION



5

10

Thus, in the broadest aspect, the present invention relates to a method of expanding and selecting disease associated T-cells, which method comprises

5

15

(a) obtaining a tissue sample from a mammal including a human being, the sample comprising disease activated T-cells, or

20

10 obtaining T-cells and antigen-presenting cell from said mammal and mixing said cells with a disease associated antigen or antigens, and

25

15 (b) culturing said tissue sample or said mixture of cells and antigen(s) in the presence of at least two factors which promote T-cell growth and optionally one or more additional compounds.

30

20 In a further aspect, the present invention relates to such continuous T-cell lines obtainable by the method.

35

35 The uses of the disease associated T-cells prepared according to the method, or the T-cell lines obtainable by the method are numerous. In particular, the T-cells and T-cell lines may be used as the active ingredient in pharmaceutical compositions and vaccines. The T-cells or  
40 T-cell lines may further be used for preparing a medicament for the treatment of various T-cell associated diseases, including diseases of inflammatory, auto-immune, allergic, neoplastic, or transplantation-related  
45 30 origin, or combinations thereof.

50

50 Furthermore, the T-cells or the T-cell lines can be used in methods for diagnosing diseases, methods for treating, alleviating or preventing diseases associated with  
35 activation of T-cells, methods of testing the effect of a

55

medicament against a T-cell associated disease, methods for the treatment, alleviation or prevention of diseases associated with T-cell activation, methods of detecting T-cell growth factors, methods of monitoring the effect of or response to treatment against T-cell associated diseases including diseases of inflammatory, auto-immune, allergic, neoplastic or transplantation-related origin or combinations thereof, and methods of identifying disease associated antigens.

Model systems for testing the effect of a medicament against T-cell associated diseases also forms part of the invention.

#### BRIEF DESCRIPTION OF THE FIGURES

Figure 1 shows schematically the T-cell vaccination principle.

Figure 2 shows schematically the establishment of a T-cell culture.

Figure 3 shows schematically the T-cell vaccination procedure.

Figure 4. Shows the number of cell population doublings, D, of three PBMC cultures grown in medium with IL-2 + IL-4 alone (left) or with allostimulation in the presence of IL-2+IL-4 (right).

Figure 5. Shows telomerase activity at 100 PD of a continuous peripheral blood activated CD4+ cell line (Act-1) cultured with IL-2 + IL-4, IL-2 or IL-4 as indicated. For comparison, telomerase activity of the leukemic cell line Se-Ax, cultured with IL-2 alone, is also shown.

Figure 6. Shows CD28 expression of the continuous peripheral blood derived CD4+ cell line Act-1 at PD 60 and 150 compared with CD4 and CD8 expression at PD 150 (Flow cytometric analysis).

Figure 7. Shows CD28 expression at different PD of the clonal T-cell line My-La, 46,XY,i(18q). Also shown is CD4 and V $\beta$ 18 expression at the different PD and CD8 expression at PD 200 (Flow cytometric analysis).

Figure 8. Shows the phenotype in the growing primary T-cell culture from which Gut<sub>1</sub>-1 is derived (Example 2) (Flow cytometric analysis).

Figure 9. Shows the phenotype of Gut<sub>R</sub>-2 in Example 2 (Flow cytometric analysis).

Figure 10. Shows the karyotype 45,XY,-20,add(1)(p36) of Gut<sub>R</sub>-2.

Figure 11. Shows the phenotype in the growing continuous T-cell culture of Gut<sub>1</sub>-1 (Example 2) (Flow cytometric analysis).

Figure 12. Shows the karyotype 47,XX<sub>1</sub>,+2,t(1;1) of Gut<sub>1</sub>-1.

Figure 13. FACS analysis of transmembrane TNF $\alpha$  in the four primary cultures. Two of the primary cultures were stimulated with super-antigen (SEA: Staphylococcus enterotoxin A). Lines indicate determination without Infliximab, and red line supplement of Infliximab to the cultures, respectively.

5

10

Figure 14. FACS analysis of transmembrane TNF $\alpha$ . Stimulation of three long term cultured cultures with super-antigen. Lines indicate determination without Infliximab and supplement of Infliximab to the cultures, respectively. (SEA: Staphylococcus enterotoxin A)

15

Figure 15. INF $\gamma$  production in primary cultures before and after supplement with Infliximab.

20

Figure 16. INF $\gamma$  and TNF $\alpha$  production in primary culture C8.3 before and after stimulation with super-antigen. (Ifx: Infliximab; SEA: Staphylococcus enterotoxin A).

25

Figure 17. INF $\gamma$  and TNF $\alpha$  production in long term cultured cultures C1x, C2x, C4.2 before and after stimulation with super-antigen. (SEA: Staphylococcus enterotoxin A).

30

Figure 18. Detection of apoptosis by Annexin FITC and propidium iodide (PI). Cells in apoptosis: FITC positive and located in lower right quadrant (LR). Cells in necrosis are double positive (FITC and PI positive and located in upper right quadrant). Negative cells located in lower left quadrant. (Ifx: Infliximab; SEA: Staphylococcus enterotoxin A, C3: complement, KL II: class II antibody (L 243 mouse anti human (Becton Dickinson))).

35

Figure 19 A-C. Coulter counter particle count with analysis of viable cells between cursor statistics.

45

Figure 20 A and B. In this figure, melanoma cells alone are shown (Fig. 20A). Furthermore, melanoma cells and cytotoxic T-cells are shown. Fig. 20B shows that melanoma cells are eliminated within 24 hours, leaving only some T-lymphocytes in the culture.

50

55

## DETAILED DESCRIPTION OF THE INVENTION

As mentioned above, the present invention is based on the recognition that certain T-cells which are associated with diseases may be expanded selectively. Such T-cells have been stimulated in vivo by a disease associated antigen or antigens. Surprisingly, such T-cells can be expanded and selected in vitro under certain conditions, whereby the T-cells escape replicative senescence and become continuous or immortal. Surprisingly, the T-cell lines maintain their antigen specificity and function during continuous culturing.

In the present invention, a cell culture system is introduced where the relevant T-cell can be expanded in practically unlimited amounts. By a quality control system (Fig. 2), it will be possible to produce T-cells which could be relevant for T-cell vaccination treatment (alternatively adjuvant treatment) in for example Crohn's disease.

In Crohn's disease the principle is based on the fact that the disease is associated with increased activity in type 1 inflammatory T-cells (IFN $\gamma$  and TNF $\alpha$ ) cells. The activity is not sufficient to activate the regulatory T-cells (IL-10 producing), but is sufficient to induce proteolytic degradation of the intestinal mucousa (active disease). In a T-cell vaccination the regulatory T-cell activity will be increased by boosting the activity by injection of attenuated activated inflammatory T-cells expanded into sufficient amounts (Fig. 3).

"Continuous" or "immortal" is intended to mean that the cells have a life-span of at least 40 PD (i.e. 1 cell becoming approximately 1 kg of cell mass), such as at least 60 PD (i.e. 1 cell becoming approximately 100 tons

5 of cell mass), preferably at least 100 PD, more  
preferably at least 150 PD, such as at least 200 PD. It  
10 is further preferred that the functional profile of the  
T-cells are not substantially altered during the  
5 continuous growth meaning that the function of the T-  
cells essentially correspond to the initial cells. In  
15 certain cases, re-activation with antigen, antibodies or  
chemical compounds may be used to activate the T-cells to  
an increased growth rate. The final aim of the invention  
10 is that an unlimited amount of specific T-cells may be  
20 produced.

The method of expanding and selecting disease associated  
T-cells of the invention comprises

15 (a) obtaining a tissue sample from a mammal including a  
human being, the sample comprising disease  
activated T-cells, or

20 obtaining T-cells and antigen-presenting cell from  
said mammal and mixing said cells with a disease  
associated antigen or antigens, and

35 (b) culturing said tissue sample or said mixture of  
25 cells and antigen(s) in the presence of at least  
40 two factors which promote T-cell growth and  
optionally one or more additional compounds.

45 "Disease associated T-cells" are intended to comprise all  
30 T-lymphocytes present at the site of disease.

50 By the term "disease activated T-cells" is meant the  
fraction of disease associated T-cells that are activated  
35 by the inflammatory process taking place at the site of  
disease.

5 In the present context, the expressions "T-cell" and "T-lymphocyte" are used interchangeably.

10 The term "disease associated antigen(s)" is intended to  
5 comprise antigen(s) (foreign or auto-antigen(s)) that initiate and maintains the inflammatory response.

15 By the term "factors which promote T-cell growth" is meant biological and/or chemical compounds, cells and the  
10 like which directly and/or indirectly stimulate T-cell growth.

20 The activated disease associated T-cells can be obtained in a tissue sample comprising such cells, which sample is  
25 taken from a mammal including a human being. Alternatively, the disease associated T-cells can be derived by obtaining T-cells and antigen presenting cells (APCs) from a mammal including a human being, and mixing  
30 such cells with a disease associated antigen or antigens. The T-cells may originate from a mammal being inflicted with a T-cell associated disease or from a healthy mammal. In particular, the tissue sample is a biopsy  
35 taken at the site of the disease. Such tissue sample is expected to further comprise antigen presenting cells as well as the antigen(s) that caused the activation of the  
25 T-cells.

40 Factors which promote T-cell growth may be selected from the group consisting of cytokines which promote T-cell  
45 growth. Examples of such cytokines are IL-2, IL-15, IL-4, IL-7, IL-9, IL-10, IL-16, and functionally similar cytokines. In particular, a combination of (1) IL-2 and/or IL-15, and (2) IL-4 and/or IL-7 and/or IL-9 may be  
50 used. In one embodiment of the present method, a combination of IL-2 and IL-4 is used. However, other T-cell growth promoting factors may also be used. Examples  
35

are combinations of ligation of the surface markers CD2, CD3 or CD28 with antibodies directed against CD2, CD3 or CD28.

By the term "functionally similar" is meant that the effect observed are comparable to the effect observed by the cytokines mentioned in the context of the present invention. These functionally similar compounds may substitute the specifically mentioned compounds in the specific process referred to.

The cytokines are preferably used in a concentration of at least 1 nM each, preferably more than 2.5 nM, more preferably than 10 nM each. The concentration of the cytokines might not be important, however, the concentration should be chosen so as to ensure growth, i.e. at least 1 nM of each. Traditionally, the concentration of a cytokine is expressed as activity in units per ml (u/ml). The person skilled in the art will readily know how to interrelate u/ml and concentration (molar, M). If nothing else is stated, it is to be assumed that 200 u/ml equals 1 nM.

The T-cells and APCs are preferably obtained from any body fluid including peripheral blood, and further from the spleen, the lymph nodes and thymus, and by spinal puncture.

The T-cells to be cultured originates preferably from a tissue sample. The tissue sample is preferably selected from a biopsy, from sputum, swaps, gastric lavage, bronchial lavage, and intestinal lavage, or body fluids such as spinal, pleural, pericardial, synovial, blood and bone marrow.



5

10

A biopsy can in principle be taken from any organ including the pancreas, the intestines, the liver, the kidneys, the lymph nodes, the breasts, and from the skin. Furthermore, peripheral blood may also be a suitable source of T-cells. Preferably the cells are taken from the organ of the disease.

15

In one embodiment of the present method, the disease associated T-cells are CD4+, CD8+ or CD4-/CD8- T-cells.

10

20

In particular, the disease associated T-cells are inflammatory, cytotoxic or regulatory T-cells.

25

Within the present context "inflammation" is defined as a general term for the local accumulation of fluid, plasma proteins, and white blood cells that is initiated by physical injury, infection, or a local immune response. This is also known as a inflammatory response. Acute inflammation is the term used to describe transient episodes, whereas chronic inflammation occurs when the infection persists or during auto-immune responses. Many different forms of inflammation are seen in different diseases. The cells that invade tissues undergoing inflammatory responses are often called inflammatory cells or an inflammatory infiltrate.

30

20

35

25

40

The majority of chronic inflammatory/auto-immune disease fall within two major groups: A type 1 chronic inflammation dominated by production of primarily IFN $\gamma$  and TNF $\alpha$  (a type 1 inflammatory cytokine profile) or a type 2 chronic inflammation dominated by production of primarily IL-4 and IL-5 (a type 2 cytokine production). Examples of type 1 chronic inflammatory/auto-immune disease are multiple sclerosis and Crohn's disease, whereas examples of type 2 chronic inflammatory diseases are asthma and long-standing severe atopic dermatitis.

45

30

50

35

55

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

As IL-4 down-regulates the production of IFN $\gamma$ , lymphocytes producing IL-4 down-regulate disease activity of a type 1 chronic inflammatory disease through an interactive cellular network. IL-4 producing T cells can thus be considered regulatory T cells in a type 1 chronic inflammatory disease, implicating that in chronic inflammatory disease type 1, the balance between cells producing type 1 cytokines like IFN $\gamma$  and TNF $\alpha$  are not sufficiently controlled by opposing regulatory T cells producing IL-4.

Conversely, as IFN $\gamma$  down-regulates production of IL-4, lymphocytes producing IFN $\gamma$  down-regulate disease activity of a type 2 chronic inflammatory disease through an interactive network. IFN $\gamma$  producing T-cells are thus considered regulatory T cells in a type 2 chronic inflammation/auto-immune reaction. In type 2 auto-immune/inflammatory disease, the type 2 cytokine production is not sufficiently controlled by opposing IFN $\gamma$  producing regulatory T-cells.

As IL-10 and TGF $\beta$  producing T-cells down-regulate chronic inflammation of both type 1 and type 2, IL-10 and TGF $\beta$  producing T-cells are for both types of chronic inflammatory diseases considered to be regulatory.

The definition of inflammatory and regulatory T-cells is thus a relative term depending on the type (type 1 or type 2) of inflammation. In type 1 chronic inflammation, the T-cells producing type 1 cytokines are considered inflammatory T-cells, and IL-4 or IL-10 and TGF $\beta$  producing T-cells are considered regulatory T-cells.

If the chronic inflammation is dominated by type 2 cytokines, the type 2 cytokine producing T-cells are

5 considered inflammatory T-cells, whereas IFN $\gamma$  and/or IL-  
10 producing T-cells are considered regulatory in this  
10 type of disease.

5 Preferably, the disease associated T-cells are associated  
with a disease of inflammatory, auto-immune, allergic,  
15 neoplastic or transplantation-related origin, or  
combinations thereof. In particular, the disease of  
inflammatory or allergic origin is a chronic inflammatory  
10 disease or a chronic allergic disease.

20 Diseases of inflammatory/auto-immune origin include  
asthma, hypersensitivity pneumonitis, interstitial lung  
disease, sarcoidosis, idiopathic pulmonary fibrosis,  
25 15 interstitial lung disease associated with Crohn's Disease  
or ulcerative colitis or Whipple's disease, interstitial  
lung disease associated with Wegeners granulomatosis or  
hypersensitivity vasculitis,

30 20 vasculitis syndromes, Hennoch-Schönleins purpura,  
Goodpastures syndrome, Wegeners granulomatosis,

35 renal diseases such as antibody mediated glomerulopathia  
as in acute glomerulonephritis, nephritis associated with  
25 systemic lupus erythematosus, nephritis associated with  
other systemic diseases such as Wegeners granulomatosis  
40 and Goodpastures syndrome and mixed connective tissue  
disease, chronic interstitial nephritis, chronic  
glomerulonephritis,

45 30 gastrointestinal diseases such as Crohn's Disease,  
Ulcerative colitis, coeliac disease, Whipple's disease,  
collagenous colitis, eosinophillic colitis, lymphatic  
50 colitis,

35

5

10

15

20

25

30

35

40

45

50

55

hepatobilliary diseases such as auto-immune hepatitis,  
alcohol induced hepatitis, periportal fibrosis, primary  
billiary cirrhosis, sclerosing colangitis,

5 disorders of the central or peripheral nervous system  
such as demyelinating disease as multiple sclerosis,  
acute disseminated encephalomyelitis, sub-acute scleros-  
ing panencephalitis,

10 skin disease such as psoriasis, atopic dermatitis,  
eczema, allergic skin disease, progressive systemic  
sclerosis (scleroderma), exfoliating dermatitis,  
pemphigus vulgaris,

15 joint diseases such as rheumatoid arthritis, ankylosing  
spondylitis, arthritis associated with psoriasis or  
inflammatory bowel disease,

muscoloskeletal diseases such as myastenia gravis,  
20 polymyositis,

endocrine diseases such as insulin dependent diabetes  
mellitus, auto-immune thyroiditis (Hashimoto),  
thyreotoxicosis, Graves,

25 diseases of the hematopoetic system such as auto-immune  
anaemia, auto-immune thrombocytopenia,

cardiovascular diseases such as cardiomyopathia,  
30 vasculitis, cardiovascular disease associated with  
systemic diseases as systemic lupus erythematosus,  
polyarthritis nodosa, rheumatoid arthritis, scleroderma,  
sarcoidosis.

35 Diseases of neoplastic origin include malignant melanoma,  
Sezary's syndrome, cutaneous T-cell lymphoma, renal cell

carcinoma, colorectal cancer, breast cancer, ovarian cancer, cancer of the uterus, prostatic cancer, hepatic carcinoma, lung cancer, and sarcoma.

Furthermore, disorders relating to transplantation may be disorders which can be treated, alleviated or prevented by use of the method of the present invention.

Chronic rejection may be related to the development of pro-inflammatory type 1 cytokine producing T-cells, and, accordingly, the expansion and selection of regulatory T-cells for adjuvant treatment in such patients may be of relevance.

In a particular embodiment of the present invention, the disease is an inflammatory bowel disease, Crohn's disease, ulcerative colitis, sclerosis, type I diabetes, rheumatoid arthritis, psoriasis, atopic dermatitis, asthma, malignant melanoma, renal carcinoma, breast cancer, lung cancer, cancer of the uterus, prostatic cancer, hepatic carcinoma, or cutaneous lymphoma.

The disease associated T-cells are preferably CD4+ (positive), CD8+, or CD4-(negative)/CD8- T-cells. The disease associated T-cells are suitably, according to the definition of inflammation, such which are inflammatory T-cells or regulatory T-cells. In one embodiment, the regulatory T-cells are cytotoxic T-cells, or CD4+ T-cells which in the case of a type 1 inflammation produce IL-4 or IL-10 and TGF $\beta$ , or in the case of a type 2 inflammation produce INF $\gamma$  or IL-10 and TGF $\beta$ . In another embodiment, the inflammatory T-cells are T-cells involved in chronic inflammatory/auto-immune diseases falling within the two major groups: A type 1 chronic inflammation dominated by production of primarily IFN $\gamma$  and TNF $\alpha$  (a type 1 inflammatory cytokine profile) or a

5 type 2 chronic inflammation dominated by production of  
primarily IL-4 and IL-5 (a type 2 cytokine production).  
10 Examples of type 1 chronic inflammatory/auto-immune  
disease are multiple sclerosis and Crohn's disease,  
5 whereas examples of type 2 chronic inflammatory diseases  
are asthma and long-standing severe atopic dermatitis.

15 In accordance with the present invention, the cells to be  
expanded and selected may optionally be cultured in the  
20 presence of at least two factors which promote T-cell  
growth and one or more additional compounds which  
preferably are such as to directly or indirectly  
interfere with T-cell growth, in particular such which  
enhance or inhibit growth of inflammatory, regulatory or  
25 cytotoxic T-cells. The function of the additional  
compound is to promote the selection and expansion of a  
desired function of the T-cells (i.e. inflammatory or  
regulatory). When such additional compound or compounds  
30 is used, it may preferably be selected from cyclosporin,  
20 GM-CSF, Prednisone, Tacrolimus, FK506, IL-10, IL-10  
antibody, TNF $\alpha$  antibody, IL-12, anti-IL-12, IL-7, anti-  
IL-7, IL-9, anti-IL-9, IL-16, caspase inhibitors, and  
35 similar compounds.

25 In another embodiment, the method comprises a selection  
procedure. Such selection procedure is described in  
40 further detail below.

45 Inflammatory cells may suitably be cells having a CD4+  
30 phenotype and a type 1 cytokine profile. The inflammatory  
T-cells are in particular cells contributing in a type 1  
inflammatory infiltrate, which cells further produce INF $\gamma$   
and TNF $\alpha$ .

50 As mentioned above, the selection is accomplished by  
35 addition of one or more additional compounds selected

5 from cyclosporine, Prednisone, Tacrolimus, FK506, GM-CSF,  
10 IL-12, IL-16, anti-IL-10, anti-TNF $\alpha$ , and functionally  
similar compounds.

5 In another aspect of the present method, the inflammatory  
T-cells are cells having a CD4+ phenotype and a type 2  
15 cytokine profile. Such inflammatory T-cells are in  
particular cells contributing in a type 2 inflammatory  
infiltrate, which cells produce IL-4 and IL-5.

10 As mentioned above, the selection is accomplished by  
addition of one or more additional compounds selected  
20 from cyclosporin, Prednisone, Tacrolimus, FK506, GM-CSF,  
IL-16, anti-IL-12, and functionally similar compounds.

25 Thus, the present invention relates to a method as  
described above, wherein the disease is mediated or  
partially mediated by type 1 or type 2 inflammatory T-  
30 cells such as chronic inflammatory bowel diseases e.g.  
Crohn's disease and ulcerative colitis, sclerosis, type I  
20 diabetes, rheumatoid arthritis, psoriasis, atopic  
dermatitis, asthma, and transplantation-related diseases.

35 In another aspect of the present method, disease  
25 associated regulatory T-cells are expanded and selected.  
Such regulatory T-cells are suitably cells having a CD4+  
40 phenotype and a type 1 cytokine profile regulating a type  
2 inflammatory disease. In particular, such cells are  
producing INF $\gamma$  and/or IL-10. Selection of such T-cells is  
30 accomplished by addition of one or more additional  
compounds selected from IL-10, IL-12 and functionally  
45 similar compounds. The invention further relates to a  
method as described above, wherein the disease is  
50 mediated or partly mediated by type 2 inflammatory T-  
35 cells, e.g. asthma or atopic dermatitis.

5 The regulatory T-cells may also be cells having a CD4+ phenotype and a type 2 cytokine profile regulating a type  
10 1 inflammatory disease. Such regulatory T-cells are cells producing IL-10 and/or IL-4. Selection of such regulatory  
5 T-cells is accomplished by addition of one or more additional compounds selected from anti-IL-12, IL-10, GM-CSF, IL-16, and functionally similar compounds. Thus, the  
15 present invention relates to a method as described above, wherein the disease is mediated or partially mediated by  
10 type 1 inflammatory T-cells e.g. chronic inflammatory bowel diseases such as Crohn's disease and ulcerative colitis, sclerosis, type 1 diabetes, rheumatoid arthritis, and psoriasis.

25 15 Furthermore, the present invention relates to a method as described above, wherein disease associated cytotoxic T-cells are expanded and selected. In particular, such cytotoxic T-cells may have a CD8+ phenotype. The  
30 cytotoxic T-cells are further preferably tumour infiltrating lymphocytes (TIL) or cells having similar properties. The CD8+ cells are often auto-immune cells that kill tumour cells. The selection of such cells are  
35 accomplished by addition of one of more additional compounds selected from GM-CSF, caspase inhibitors such as Z-VAD,  $\alpha$ -CD95, IL-10, IL-12, IL-16, and functionally  
25 similar compounds. The present invention relates to a method as described above, wherein the disease is of neoplastic origin such as malignant melanoma, renal carcinoma, breast cancer, lung cancer, cancer of the  
40 uterus, prostatic cancer, hepatic carcinoma, and cutaneous lymphoma.

50 In a further aspect, the present invention relates to continuous T-cell lines obtainable by the methods as  
35 defined above and claimed herein. In particular, the T-



5

cell line is such, wherein the T-cells are inflammatory T-cells, regulatory T-cells or cytotoxic T-cells.

10

As demonstrated in Example 1, the antigen specific T-cells overgrow T-lymphocytes not having the desired specificity. It should be noted that in the examples shown, the T-cells with the shortest PD-time (i.e. the fastest growing T-cells) would preferentially be expanded. In general, it is not to be expected that T-lymphocytes with a desired specificity, avidity, growth potential, phenotype and function preferentially expand over T-cells with other antigen specificities. However, the realisation that antigen specific T-cells can be obtained in an unlimited number implies that appropriate selection procedures will be able to establish T-lymphocyte cell lines with the desired specificity, avidity, growth potential, phenotype and function.

15

20

25

30

As discussed above, in many chronic diseases, the natural balance between inflammatory and regulatory T-cells has been disrupted and cannot find it's way back in balance. For each such disease, it would be possible to select for and expand either inflammatory T-cells or regulatory T-cells. Because of the in vivo activation of the T-cells, the selected and expanded T-cells are antigen-specific, and thus disease-specific. Dependent on the desired route of treatment, the selection of inflammatory T-cells (T-cell vaccination) or regulatory T-cells (adjuvant treatment) may be directed.

35

40

45

Selection for antigen specific T-cell growth is initiated by antigen presentation. In case of a biopsy harbouring disease associated lymphocytes, it is assumed that the biopsy initially, besides the disease associated T-lymphocytes, also contains antigen and antigen presenting cells. Upon expansion of T-lymphocytes, the initial

50

55

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

activation by antigen may not be sufficient for continuous T-lymphocyte growth, and in vitro activation of the desired T-lymphocytes may be necessary. In vitro activation requires access to autologous or HLA matched antigen presenting cells. These can be obtained from a blood sample, as so-called mononuclear cells. Furthermore, powerful antigen presenting cells (dendritic cells) can be obtained from mononuclear cells by culturing plastic adherent mononuclear cells in a medium supplemented with granulocyte-macrophage colony stimulating factor (GM-CSF) and IL-4 (both at a concentration above 1000 u/ml). Dendritic cells will develop within 8-20 days.

15  
20  
25  
30  
35  
40  
45  
50  
55

Having obtained antigen presenting dendritic cells and disease associated T-lymphocytes, a preferential growth advantage of antigen specific T-lymphocytes is to be expected by mixing antigen, dendritic cells and disease associated T-lymphocytes, or peripheral autologous mononuclear cells, as a source of T-lymphocytes in case a biopsy from the diseased organ is not available. The medium should at least contain two factors promoting T-cell growth and an additional factor, the latter to secure transient growth and differentiation of dendritic cells in cases dendritic cells are necessary. A combination of such factors could be the following cytokines: IL-2, IL-4 and GM-CSF. Furthermore, human serum is preferred in order to minimise the autologous mixed leukocyte reaction.

30  
35  
40  
45  
50  
55

However, antigen activation of T-lymphocytes may lead to proliferation as well as to activation induced cell death (AICD). The balance between proliferation and cell death determines the growth rate (positive or negative) of a cell culture. In order to down-regulate AICD, inhibitors of AICD can be included in the growth medium. Examples of

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

such inhibitors are caspase inhibitors (like Z-VAD) and certain antibodies with reactivity to CD95 (Fas) that prevents Fas-FasL induced cell death. In addition antigen activation of T-lymphocytes may lead to development of T-cells not having the desired phenotype and/or function, implying how further selection and/or counter-selection procedures can be carried out in order to obtain continuous T-lymphocyte cell lines with the desired properties (cf. below).

10  
20  
25  
30  
35  
40  
45  
50  
55

As the cell culture system promotes the expansion of the fastest growing T-cell clones, bystander cells not having the desired specificity may overgrow the ones having the wanted specificity, reactivity, phenotype and function. As T-lymphocyte growth in general is dependent on IL-2 as well as IL-4, growth cessation may be obtained by withdrawal of one or both of these cytokines. Specific antigen activation of growth arrested T-lymphocyte cell lines is expected to favour proliferation of antigen specific T-cells in a medium with at least two cytokines.

35  
40  
45  
50  
55

It is important to monitor activation of the T-lymphocytes, as this shows whether the antigen activation is successful, and gives additional information concerning selection/counter-selection of the desired T-lymphocyte sub-population.

30  
35  
40  
45  
50  
55

Several assays are available to monitor T-cell activation. Activation markers induced on the surface of the T-lymphocytes by antigen activation, such as CD25, CD69 and membrane bound TNF $\alpha$  may be used to measure the degree of activation, and may also be used by immuno-separation techniques to select for antigen activated T-lymphocytes. Similarly, differentiation markers such as CD4 and CD8 may be used by immuno-separation techniques to select for T-cells with the appropriate phenotype.

5

10

15

Furthermore, selection of sub-populations of T-lymphocytes expressing particular  $V_\alpha$  and  $V_\beta$  subfamilies of the T-cell receptor complex may be very useful. Importantly, if the antigenic peptides bound to the major histocompatibility complex (MHC) are known, peptide-MHC tetramers can be used to immuno-select T-lymphocytes with the desired specificity and avidity.

20

25

30

35

40

45

50

55

Effector functions like cytokine production and cell killing gives information regarding the strength of the antigen activation. However, antigen activation of a given sub-population may activate the immunological network given rise to the outgrowth of regulatory T-cells capable of down-regulating the desired sub-population of T-lymphocytes. As an example of this phenomenon, it is believed that in Crohn's Disease the balance between inflammatory T-lymphocytes producing IFN $\gamma$  and TNF $\alpha$  and regulatory T-cells mainly producing IL-10 has shifted towards the inflammatory T-lymphocytes. However, a powerful activation and expansion of clonal inflammatory T-lymphocytes is expected to be followed by activation and expansion of regulatory T-lymphocytes, which participate in a down regulation of the inflammatory response. In this way the T-cell vaccination with activated and attenuated inflammatory T-lymphocytes results in a down regulation of the disease related elevated level of inflammatory T-lymphocytes. In this case in order to minimise the establishment of regulatory T lymphocytes, addition of cyclosporin A or glucocorticoids, that partially inhibits the inflammatory response, may be useful. In addition, as IL-10 is of importance for establishment of regulatory CD4 $^+$  T-cells, neutralising antibodies to IL-10 may be added to the medium. Conversely, if regulatory lymphocytes are to be established, inflammatory T-lymphocytes should be highly

5 activated, and/or IL-10 added to the medium containing at  
10 least one additional cytokine.

10 Apart from antigen activation, other non-specific methods  
5 are available that promote T-cell growth, and if combined  
with appropriate selection procedures as outlined above,  
15 may enhance T-lymphocyte growth, in cases where the cell  
population doubling time is considered too long. Such  
methods include activation by super-antigen pulsed  
10 antigen presenting cells, activation by mitogens (like  
20 PHA and jacalin) in the presence of feeder cells or  
antigen presenting cells, activation by antibodies  
against CD2, CD3 and CD28, activation by ionomycin and  
phorbol ester and in case of cross-reactivity with  
25 15 alloantigen, allostimulation with appropriate allogenic  
cells with or without autologous dendritic cells (the  
latter possibility in order to obtain cross-priming).  
AICD can in all the cases mentioned above be blocked by  
30 caspase inhibitors.

20 The principles outlined above are also applicable if  
35 cloned T-cells with a given specificity are available.

25 The disease determines the subtypes of T-cells which  
could be relevant as a treatment principle. In auto-  
40 immune disease T-cell vaccination with a disease antigen,  
associated pro-inflammatory type 1 cytokine profile (IFN $\gamma$   
and TNF $\alpha$ ) T-cell line could be relevant. If it is not  
possible to select the disease associated antigen  
45 30 reactive pro-inflammatory T-cell line, it may be possible  
to select a regulatory T-cell line with a type 2 cytokine  
profile (IL-4/IL-10) which, in an analogous fashion, can  
be used as a immunoadjuvant therapy against the disease  
50 associated inflammatory T-cells.

35

5 In order to select for the desired type of T-cell  
immunological selection principles or additional  
10 compounds can be used as described above.

5 In the following, a selection of important diseases in  
relation to the present invention is discussed.

15 Crohn's Disease

The chronic inflammatory disease Morbus Crohn (Mb. Crohn,  
10 Crohn's Disease) is a relatively frequently occurring  
20 disease, the prevalence being 55 per 100000 individuals.  
The incidence has during the last 20 years been  
increasing by 8-9 new cases per 100000 individuals per  
year. Diagnosis and treatment of Crohn's disease are  
25 therefore a major task for specialised medical  
gastroenterologic hospitals.

In the past, the treatment of Crohn's disease has been  
30 based on an inhibition or modulation of the immune system  
20 by means of i.a. azathioprine and cyclosporin. The  
results obtained by this treatment have been varying, and  
a way of dividing the disease into subgroups may be  
35 needed in order to successfully treat the disease by  
immune modulation.

25 Recent research has rendered it possible that Morbus  
40 Crohn is a multi-factorial auto-immune disease. It has  
been suggested that the normal tolerance of the immune  
system against the microbial flora in the intestines are  
30 broken (ref. 6). The chronic immune reactivity against  
45 the bacterial flora seems to be mediated by T-lymphocytes  
producing INF $\gamma$  and TNF $\alpha$ . The constant presence of these  
cytokines in increased amounts contributes to the  
50 destruction of tissue (an auto-immune reaction) which  
35 take place in the inflamed intestine. The treatment of  
Crohn's disease has accordingly during phase 1 and 2

clinical studies been focused on modulation of the T-cell-mediated immune response by use of IL-10, CD4 antibodies and antibodies against TNF $\alpha$  (refs. 7, 8, 9).

As mentioned above, Crohn's Disease is believed to be a multifactorial disease associated with pro-inflammatory IFN $\gamma$  and TNF $\alpha$  producing T-cells in the intestinal mucosa. Basically the balance between the pro-inflammatory T-cells and regulatory T-cells is dysregulated resulting in increased production of the pro-inflammatory cytokines. The fundamentals for T-cell vaccination is based on these observations. The pro-inflammatory immune response is activated by different disease relevant antigens. Nevertheless, the activation level in vivo/in situ is not sufficient to activate the regulatory immune response. To stimulate the in vivo regulatory immune response, activated pro-inflammatory T-cells are selected, cultured, activated and attenuated and administered to the patient.

The culture system of the invention selects for the T-cell line with the shortest PD time as shown in example 2. In this case the pro-inflammatory cytokine producing CD4 $^{+}$  T-cell line expands from the gut biopsy on behalf of the regulatory T-cells. In order to avoid the propagation of IL-10 producing regulatory T-cells, which suppress the growth of pro-inflammatory T-cells, selection procedures, as described above can be used. Cyclosporine suppresses the production of IFN $\gamma$  and TNF $\alpha$  of the in vivo antigen stimulated pro-inflammatory culture (ref. 6). In cultures where cyclosporine is used as a supplement to at least two cytokines, the development of regulatory T-cells is suppressed. Regulatory T-cells are dependent on the presence of IL-10 or/and TGF $\beta$ , and in order to establish pro-inflammatory T-cells from intestinal biopsy specimen the selection of pro-inflammatory T-cells may be

5 facilitated by the addition of IL-10 antibody to early  
cultures. Of course combination of antibody to IL-10 and  
10 cyclosporine may also be used.

5 If the established culture is not sufficiently growing,  
it can be stimulated with autologous relevant antigen,  
15 either intestinal sonicated bacterial material presented  
by antigen presenting cells (dendritic cells developed  
from peripheral blood), or by auto-presentation of super-  
20 antigen in accordance with the Examples below, or  
presented with pulsed APCs. To avoid activation induced  
apoptosis,  $\alpha$ CD95 or Z-VAD could be used concomitantly in  
the culture medium.

25 15 The development of dendritic cells is dependent on the  
presence of GM-CSF and IL-4 (ref. 7). When a sufficient  
amount of dendritic cells are available ( $10^7$ )  $10^6$   $\gamma$ -  
30 irradiated dendritic cells incubated with sonicated  
bacterial material or super-antigen are mixed with the  
20 desired culture in a 1:1 relationship. After 24 hours,  
positive selection may be performed by usage of either  
CD69-Ab, CD25-Ab, FAB210 (transmembrane TNF $\alpha$  antibody) or  
35 Infliximab (chimeric TNF $\alpha$  antibody with high avidity for  
transmembrane TNF $\alpha$ ).

25 In patients with severe disease, the activity level of  
40 the inflammatory T-cell is pronounced and if the  
development of IL-10 producing T-cells is avoided,  
cultures relevant for T-cell vaccination emerges. In  
45 30 cases where the in vivo antigen activation elicits a  
regulatory T-cell response, selection of pro-inflammatory  
T-cells by antibodies against the activation markers  
CD69, CD25 or transmembrane TNF $\alpha$  is an option in the  
50 early phase of the culture. Usually regulatory cells do  
35 not establish until two to three weeks after the  
establishment of the culture and the time related



dynamics in the culture can be used in the selection process. Magnetic beads coupled to the relevant activation marker antibody may for instance be used.

The expression of surface activation markers and proliferation can also be non-specific augmented by CD3-Ab, CD2-Ab, CD28-Ab. Positive selection after stimulation is performed as described above.

In some cultures the growth of CD4+ cells could be inhibited by CD8+ cells. The CD8+ cells can be removed by negative selection.

When  $10^3$ - $10^{10}$  cells with the relevant phenotype (CD4+, CD45RO+, CD25+, (Act-1)+, CD69+, Transmembrane - TNF+) and function (IFN $\gamma$  and TNF $\alpha$  production) are available, the cells may advantageously be activated and attenuated by  $\gamma$ -irradiation prior to administration to the patient, for example in the form of an injection subcutaneously in the forearm.

Selection of regulatory T-cells for adjuvant therapy in patients with Crohn's Disease can be achieved by allostimulation with the allogenic T-cell line Se-Ax (cf. the Examples). It is assumed that the pro-inflammatory T-cell recognises the allogenic Se-Ax (an IL-10 producing T-cell line from a patient with Sezary's syndrome). Hereby a pro-inflammatory response inducing secretion of type 1 cytokines stimulate the development of regulatory T-cells (because Se-Ax also produces IL-10 needed to generate CD4+ regulatory lymphocytes). Regulatory T-cells can also be induced by the addition of IL-10 to the culture media also in combination with TGF $\beta$  (ref. 10). The autologous regulatory IL-10 producing T-lymphocytes may be used as intravenous adjuvant immunological therapy in patients with active Crohn's Disease.

Different patients with Crohn's Disease may share common peptides in the variable region of the  $\beta$ -chain of the T-lymphocyte receptor site essential for the development of the type 1 response. In this case peptide libraries from the T-cell receptor  $V_{\beta}$ -chain could be used as a vaccine in Crohn's Disease.

#### Asthma

Asthma is related to type 2 cytokine producing (IL-4, IL-5, IL-3 and GM-CSF) T-lymphocytes in the bronchial epithelium. The cytokines mobilise and activate eosinophils for subsequent mucosal tissue injury. The same relationship is related to atopic dermatitis. In the bronchial epithelium in patients with asthma, stimulation with house dust mite (HDM) is associated with a type 2 cytokine response with production of IL-4, IL-5 and IL-10. In normal individuals, stimulation of respiratory epithelial T-lymphocytes with HDM elicits a type 1 cytokine response predominated by the production of IFN $\gamma$ .

Patients with asthma could also be subjects for T-cell vaccination with attenuated type 2 cytokine producing CD4 $^{+}$  T-cells in order to obtain IL-10 producing cells or a type 1 cytokine response reducing disease activity.

The relevant T-lymphocytes could be obtained by either bronchial biopsies or bronchioalveolar lavage and cultured in a medium supplemented with at least two cytokines, and GM-CSF. It would be relevant to use GM-CSF because dendritic cells are very abundant in the respiratory epithelium.

Dendritic cells could also be cultured according to the methods mentioned above, e.g. from peripheral blood cells.

5  
10 In asthma, it may be relevant to stimulate blood mononuclear cells with known antigens. It has previously been demonstrated, that in patients with severe asthma, 5 CD4 and CD8 enriched peripheral blood expresses spontaneously increased amounts of mRNA for the type 2 cytokines localised to CD4 but not CD8 cells (ref. 11). 15 These CD4+ cells could be stimulated with antigen presented by dendritic cells, developed as mentioned previously, in a medium supplemented with high levels of 20 IL-2 and IL-4 as described by Kaltoft 1998 (ref. 12).

25 The T-lymphocytes obtained by culture should be described concerning function, avidity (known antigen), and phenotype. Selection procedures as described previously 30 can be used. Cyclosporine has been used in the treatment of asthma. In these patients a down-regulation of the IL-5 response is important, probably because of inhibited IL-5 gene transcription by cyclosporine (and FK506), but 35 also because of a general down-regulation of calcium dependent transcription of cytokine mRNA (ref. 13). In order to eliminate a type 1 T-cell overgrowth in the established cultures, cyclosporine may be added to the cultures.

40 25 In many cases of asthma, the antigen is known (HDM or T-cell reactive peptides in asthma ICP1 and ICP2 epitopes known in cat allergy (synthesised from the cat allergen Fel d1)). In order to stimulate growth, HDM, ICP1, ICP2 45 30 or other relevant antigens can be used presented by dendritic cells. In asthma, co-stimulation of the T-cells with dendritic cells via CD28 could be combined with CTLA4-Ig fusion protein because when dendritic cells 50 35 ligate with CTLA 4 on T-cells, it has been associated with apoptosis.

5

Multiple Sclerosis

10

The disease is associated with auto-immune CD4+ T-cells reactive against the myelin, inducing secretion of inflammatory type 1 cytokines in the diseased neural tissue. In multiple sclerosis, T-cell vaccination has been investigated, but so far, it has not been possible to obtain sufficient amounts of activated T-cells with the desired phenotype and cytokine profile. In previous vaccination attempts, the disease associate phenotype, specificity and function of the T-lymphocytes have not been secured (refs. 14, 15).

15

20

25

The culture and selection procedure which could be relevant in these patients are similar to the methods described for Crohn's Disease.

Relevant in vivo disease associated, in vivo antigen activated T-cells could be obtained by spinal puncture. This material could after centrifugation, be propagated in a medium containing at least two cytokines. If growth is not sufficient the T-cells could be activated with myelin as described above.

30

20

35

If sufficient amounts of T-cells cannot be obtained from spinal puncture, peripheral blood mononuclear cells could be separated and isolated by a Ficoll-Isopaque gradient and stimulated by APC presenting myelin.

40

Continuous CD4+ T-cells with reactivity against myelin and a type 1 cytokine production will after activation and attenuation be ready for T-cell vaccination.

45

30

Cancer

For the treatment of cancer, the present invention is believed to be of special interest. Most cancers are associated with tumour infiltrating lymphocytes (TIL),

50

35

55

5  
10  
5 and these TIL's are known to have killer cell activity against the tumour cells. Examples of cancers where this phenomenon are well documented include melanoma, colorectal cancer, renal cell carcinoma, breast cancer and sarcoma.

15  
10 Although TIL's have anti-tumour activity, the main problem for efficient treatment of cancer with TIL's have so far been that it has not been possible to grow TIL's in sufficient quantities.

20  
25  
30  
20 TIL's have so far been cultured to approximately  $10^{11}$  cells (ref. 16) corresponding to approximately 100-300 grams of cells and this quantity has in most cases not been sufficient to combat large tumour masses (in the order of some kg) also partly because not all the cultured TIL's after long term culture do not have the desired specificity against the tumour cells (ref. 16). The present invention overcomes these limitations. So far it has been shown that two continuous CD8+ T-cell lines with specificity and killer cell activity against autologous tumour cells have been established from patients with mycosis fungoides and Sezary's syndrome, respectively.

25  
40  
45  
30 Examples of cancerous diseases which could be treated with the T-cell lines or T-cells prepared according to the present invention include malignant melanoma, renal carcinoma, breast cancer, lung cancer, cancer of the uterus, prostatic cancer, cutaneous lymphoma and hepatic carcinoma.

50  
35 As most tumour associated antigens are relatively few (because most tumour associated antigens are self-antigens), the present invention as outlined herein may be used not only to treat the patient from which the

lymphocytes derived, but also offers the possibility of treating different but HLA matched patients with these established continuous T-cell lines.

For example in the case of metastatic malignant melanoma, HLA-typing may be performed on peripheral blood cells. If the result of this typing shows that the patient for example expresses HLA-A2 (HLA 0201), the immunogenic melanoma associated peptides restricted by this HLA allele are known to derive from at least the following proteins: Tyrosinase, Melan-A/Mart-1 and gp100. The amino acid sequence of the HLA-A2 binding melanoma associated peptides is for tyrosinase MLLAVLYCL, for Melan-A/Mart-1 AAGIGILTV, and for gp100 KTWGQYWQV. Peptide-MHC tetramers from these melanoma associated peptides can then be used to determine whether the patient has circulating CD8+ T-lymphocytes with specificity to the peptides. Such CD8+ positive cells are also expected to be present in a larger fraction of the outgrowing biopsy derived T-lymphocytes, and the peptide-MHC tetramer technique can thus be used to enumerate and select for melanoma antigen specific T-cells with different avidity among outgrowing biopsy derived T-lymphocytes. Furthermore, immunochemistry of tumour biopsy material can confirm and supplement the data obtained by the peptide-MHC tetramer technique.

The outgrowing T-lymphocytes are in general of oligoclonal origin and consist of both CD4+ and CD8+ T-lymphocytes. Contained within the latter population are the presumed auto-immune effector cells (killer cells), while contained within the former population are CD4+ cells mediating help in generating CD8+ effector cells.

Peripheral blood derived dendritic cells can be pulsed with melanoma associated peptides and used to expand

5  
10 melanoma associated peptide-MHC tetramere selected CD8+ cells in the medium supplemented with IL-2, IL-4, GM-CSF and Z-VAD or functionally similar combinations of growth and selection factors in the presence of  $\gamma$ -irradiated  
15 outgrowing T-cells as feeder and helper cells. The desired CD8+ tumour specific lymphocytes may then be expanded according to the procedures aiming at expanding T-lymphocytes in an unlimited number. The specificity and function of the T-lymphocyte cell lines can be confirmed  
20 by killing and cytokine production of HLA-matched tumour cells presenting the melanoma associated peptides in question.

25 If the melanoma associated peptides are not known, but melanoma cells or melanoma cell lysate are available an alternative approach can be employed. Dendritic cells and outgrowing lymphocytes are mixed for some time in a medium containing IL-2, IL-4, GM-CSF and Z-VAD or  
30 functionally similar combinations of growth and selection factors. Later tumour cells or tumour cell lysate are added and following expansion, appropriate selection procedures should select for CD8+ cells with tumour cell reactivity. It should be noted that continuous T-cell  
35 lines are often oligoclonal for more than 100 PD, implying that continuous CD8+ tumour specific T-lymphocyte cell lines may react with several melanoma  
40 associated antigens, thus minimising the risk of tumour escape.

45 30 Selection for melanoma specific CD8+ cells may also be obtained by mixing outgrowing T-lymphocytes with tumour cells in a medium with IL-2, IL-4 and Z-VAD or functionally similar combinations of growth and selection  
50 factors, because the tumour cells (target cells) acts as antigen presenting cells by directly presenting tumour associated peptides to CD8+ T-lymphocytes.  
55

5  
10  
15  
20  
25  
30  
20  
35  
40  
45  
50  
55

When continuous CD8+ T-cell lines are available, these cell lines can be used to treat HLA-matched melanoma patients tumour associated antigens recognised by the continuous CD8+ cell lines (including of course the patient from which the continuous cell lines derive), in particular patients with metastatic malignant melanoma. In the case described above melanoma patients with HLA-A2, an allele which more than 40% of Caucasian melanoma patients carry. Patients with metastatic malignant melanoma have a very poor prognosis with a median survival time of only 7.5 months. Accordingly it is desirable to have access to treatment options that can work fast like pre-made continuous HLA-matched tumour specific CD8+ cell lines. As the vast majority of HLA-matched melanoma patients express the same tumour associated antigens, it may be possible to establish a T-lymphocyte cell bank that optimally will fit every patient with malignant melanoma regarding tumour cell killing and HLA-match for non-presenting HLA-alleles.

35  
40  
45  
50  
55

The tumour specific CD8+ T-lymphocytes may be  $\gamma$ -irradiated in order to ensure that the cells cannot divide further and infused to the patient in combination with an established IL-2 therapy protocol. Before administration, e.g. infusion, the T-lymphocytes can be incubated with the caspase inhibitor Z-VAD, in order to reduce AICD, or Z-VAD may be given during the administration. Like other TIL's, the continuous CD8+ cell lines are expected to home to the tumour bed, thereby initiating a massive tumour cell destruction followed by cytokine production located at the tumour sites. Besides killing of the tumour, AICD is expected to lead to a fast elimination of the administered lymphocytes, which in general should be sensitive to Fas-FasL killing in order not gain access to immune



5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

privileged sites such as the eyes and the testis. Due to the elimination of lymphocytes, large quantities are obviously needed to combat large tumour masses. As soluble melanoma associated peptide HLA complexes are released during melanoma cell killing, such complexes interfere with the interaction between CD8+ cells and melanoma cells. Thus, it may be necessary to remove such complexes from the blood stream during treatment for instance by an immuno-magnetic separation technique. The presence of soluble melanoma peptide HLA complexes can however serve as a marker for the effectiveness of tumour eradication. Furthermore, when allogenic cytotoxic cells differ from the patient's HLA-type, this may be used to follow the number and fate of the infused lymphocytes.

15  
20  
25  
30  
35  
40  
45  
50  
55

It is expected that the administered attenuated, for example infused, e.g.  $\gamma$ -irradiated, CD8+ lymphocytes are capable of killing, if not all, then the vast majority of tumour cells. Furthermore the inflammation generated by the administered CD8+ cells (perhaps also with the addition of administered helper CD4+ cells) will activate autologous resident but inactive melanoma specific pre-cytotoxic T-cells to killer cells, in part due to the expansion/maturation of dendritic cells that are activated by the production of GM-CSF and TNF $\alpha$  during melanoma cell killing.

In combination, the above effect mechanisms are expected to eradicate all tumour cells.

#### 30 35 40 45 50 55

Pharmaceutical compositions

The present invention also relates to pharmaceutical compositions comprising activated disease associated T-cells prepared according to the methods described herein, or comprising one or more T-cell lines as described

herein, optionally comprising one or more pharmaceutically acceptable drugs and/or excipients.

The T-cells to be used in the composition are preferably inflammatory T-cells, regulatory T-cells, or cytotoxic T-cells.

In one embodiment, the composition comprises T-cells or one or more T-cell lines which have been re-activated in the presence of one or more antigens. Such antigens may preferably be disease associated antigen(s), alloantigen(s), or super-antigen(s). Examples of super-antigens are SEA, SEB, SEC, SED, SEE, TSST, Streptococcus pyogenes enterotoxin A, B and C, and Mycoplasma arthritidis antigen. Disease associated antigen(s) can be added in the event the antigen is known. Alternatively, re-activation may be carried out with a tissue sample or another sample expected to comprise the disease associated antigen.

The T-cells are preferably attenuated prior to administration in order to ensure that the cells are not able to divide further. Such attenuation may suitably be accomplished by x-ray or UV radiation or by addition of cell poisons.

The suitable amount of the T-cells of the invention to be administered depends on several factors, i.a. the disease or condition to be treated, alleviated or prevented, and further on the age, weight and state of the subject to be treated. The skilled person art will readily know how to establish the optimum dose.

The administration may be as single doses or as several doses per day. In certain cases, administration only once may be sufficient. In general, several doses should be

5 given such as once for a period of for examples a day for  
a week or for months, or repeated administration once  
10 every week, every second week, etc.

5 The amount of the T-cells of the invention depends on  
patient, on the route of administration, and the severity  
15 of the disease or condition to be treated. In general,  
 $10^8$ - $10^{12}$  cells may be suitable for each dose.

10 The pharmaceutical composition is conveniently  
administered parenterally, by injection either  
20 subcutaneously, intramuscularly, intravenously or by  
infusion.

25 For  $V_{\beta}$  disease specific peptides, injectables may be in  
the form of liquid suspensions or solutions, solid forms  
suitable for solubilisation or suspension in liquid prior  
30 to injection. The pharmaceutical composition may also be  
emulsified. Additional modes of administration may in  
20 certain cases be suitable such as e.g. oral formulations.

35 The pharmaceutical composition may also be mixed with  
suitable excipients such as e.g. water, saline, dextrose,  
glycerol, ethanol or combinations thereof. In addition,  
25 the composition may contain auxiliary substances such as  
40 wetting agent, emulsifying agents, colouring substances,  
preserving agents, or pH buffering agents.

#### Vaccines

45 30 T-cell vaccination seems to be an attractive treatment of  
various diseases including auto-immune diseases and  
cancer. However, in practice, T-cell vaccination has not  
50 been a realistic option since auto-reactive T-cells as  
35 other humane T-lymphocytes are believed to have a limited  
dividing capacity in vitro.

5

10

One problem is that it has not been possible to obtain a sufficient number of cells to perform vaccination. By the present invention, an unlimited number of cells is available, thus, making T-cell vaccination possible.

15

20

Accordingly, in another aspect, the present invention relates to vaccines comprising activated disease associated inflammatory T-cells prepared in accordance with the methods described herein, or one or more T-cell lines as described herein.

25

30

In one embodiment of the vaccine, the T-cells have been re-activated in the presence of one or more antigens. Representative examples of such antigens are disease associated antigen(s), alloantigen(s), or super-antigen(s). Examples of super-antigens are SEA, SEB, SEC, SED, SEE, Streptococcus pyogenes enterotoxin A and B, and Mycoplasma arthritidis antigen.

20

35

In a preferred embodiment of the vaccine, the T-cells have been attenuated. Such attenuation may suitably be accomplished by  $\gamma$ - or UV-radiation, or by addition of cell poisons.

25

40

Disease associated antigen(s) can be added in the event the antigen is known. Alternatively, re-activation may be carried out with a tissue sample or another sample expected to comprise the disease associated antigen.

45

30

50

The T-cells are preferably attenuated prior to administration in order to ensure that the cells are not able to divide further. Such attenuation may suitably be accomplished by x-ray or UV radiation or by addition of cell poisons.

35

55

5  
10 The suitable amount of the T-cells of the invention to be administered depends on several factors, i.a. the disease or condition to be treated, alleviated or prevented, and further on the age, weight and state of the subject to be  
5 treated. The skilled person art will readily know how to establish the optimum dose.

15  
20 The administration may be as single doses or as several doses per day. In certain cases, administration only once may be sufficient. In general, several doses should be given such as once for a period of for examples a day for a week or for months, or repeated administration once every week, every second week, etc.

25  
30 15 The amount of the T-cells of the invention depends on patient, on the route of administration, and the severity of the disease or condition to be treated. In general,  $10^8$ - $10^{12}$  cells may be suitable for each dose.

35  
40 20 The pharmaceutical composition is conveniently administrated parenterally, by injection either subcutaneously, intramuscularly, intravenously or by infusion.

45  
50 25 For  $V_p$  disease specific peptides, injectables may be in the form of liquid suspensions or solutions, solid forms suitable for solubilisation or suspension in liquid prior to injection. The pharmaceutical composition may also be emulsified. Additional modes of administration may in  
30 certain cases be suitable such as e.g. oral formulations.

55  
35 The pharmaceutical composition may also be mixed with suitable excipients such as e.g. water, saline, dextrose, glycerol, ethanol or combinations thereof. In addition, the composition may contain auxiliary substances such as

wetting agent, emulsifying agents, colouring substances, preserving agents, or pH buffering agents.

Prior to vaccination with the vaccine of the present invention, or treatment with the pharmaceutical composition of the present invention, the phenotype for T-lymphocyte receptors TCR- $\alpha$ , $\beta$  and TCR- $\gamma$ , $\delta$  of the cell culture may be determined e.g. by flow cytometry. Likewise, the HLA-DR, CD3, CD4, CD8, CD11, CD18, CD23, CD28, CD45RO, CD54, HML-1 CD11a and clonality characteristics may be determined, providing important information.

Furthermore, the cytokine profile may be determined. The following cytokines may be determined: INF $\gamma$ , IL-10, TNF $\alpha$ , IL-12, IL-2, IL-4 and TGF $\beta$ . Also, extended HLA class I and II as well as status regarding Hepatitis A, B, and C, and HIV, CMV, EBV and HTLV-I should be determined for both the subject and the cell lines. Also, intracellular amount of NF $\kappa$ B and JAK/STAT pathway may be monitored.

#### Uses of the T-cells or T-cell lines of the present invention

Furthermore, the use of the T-cell lines and T-cells as described and claimed herein of a medicament for the treatment of a T-cell associated disease also forms part of the present invention. In particular, the medicament is used for treating, alleviating or preventing diseases of inflammatory, auto-immune or neoplastic origin, or combinations thereof. Examples of such diseases are given above. In particular, the medicament may be for treating, alleviating, or preventing inflammatory bowel disease, Crohn's colitis, sclerosis, type I diabetes, rheumatoid arthritis, psoriasis, atopic dermatitis, malign melanoma,

5  
10 renal carcinoma, breast cancer, cutaneous lymphoma, or the like.

10  
5 In addition to the content of T-cells, the compositions or vaccines may contain drugs for use in a conventional treatment of the particular disease, or drugs for the treatment or prevention of side effects in connection with the disease or treatment of the disease. Such drugs should readily be known to the practitioner (doctors etc.). Examples are 5-aminosalicylic acid, azathioprin, Prednisone, budesonide.

#### Diagnosis

25  
15 In yet another aspect, the present invention relates to a method for the diagnosis of a disease in a mammal, which method comprises

- 30  
20 (a) obtaining a tissue sample from a mammal including a human being, the sample comprising activated T-cells, antigen presenting cells, and antigen(s),
- 35  
25 (b) culturing said tissue sample or said activated T-cells in the presence of two or more T-cell growth factors and optionally one or more additional compound,
- 40  
45 (c) observing the presence of disease associated T-cells, and relating the presence of these T-cells to a disease.

50  
35 In one embodiment of the diagnostic method, the disease is related to the disease associated T-cells by determining the kind or phenotype of the activated T-cells and/or their state of activation.

5  
10 In another embodiment of the diagnostic method, the cytokine profile of the T-cells is determined. Thereby, the activated T-cells are determined, and thus, the disease.

15 Methods for the treatment, alleviation or prevention of diseases associated with activation of T-cells

20 In a special aspect, the present invention relates to a method for the treatment, alleviation or prevention of a disease associated with an activation of T-cells in a subject comprising administering to the subject one or more T-cell lines, T-cells, a composition or a vaccine as defined and claimed herein.

25 Such method comprises

30 (a) obtaining a tissue sample from a mammal including a human being, the sample comprising disease activated T-cells, or

35 obtaining T-cells and antigen-presenting cell from said mammal and mixing said cells with a disease associated antigen or antigens, and

40 (b) culturing said tissue sample or said mixture of cells and antigen(s) in the presence of at least two factors promoting T-cell growth and optionally one or more additional compound.

45 Factors that promote T-cell growth are given above and include cytokines that promote T-cell growth. Examples are IL-2, IL-4, IL-7, IL-9, IL-10, IL-15, IL-16, and functionally similar compounds. In particular embodiment, a combination of IL-2 and/or IL-15 and IL-4 and/or IL-7



5 is used, preferably a combination of IL-2 and IL-4. The  
concentration of the cytokines may preferably be at least  
10 1 nM, more preferably more than 2.5 nM, and most  
preferably more than 10 nM.

5 The method of expanding and selecting the disease  
15 associated T-cells are described in greater detail above.

20 The sample to be cultured may be a tissue sample or  
another sample as defined above. The sample from which  
the T-cells are expanded may in one embodiment be a  
tissue sample collected from the patient to be treated,  
and in another embodiment a tissue sample collected from  
a patient different to the patient to be treated.  
25 Furthermore, the HLA restriction of the T-cells and in  
the patient to be treated may be determined.

30 Diseases to be alleviated, prevented or treated are in  
particular those described above.

20 Furthermore, the invention relates to a method for the  
treatment, alleviation or prevention of a disease  
35 associated with an activation of T-cells in a subject  
comprising administering a medicament as identified  
25 according to the method identified as being effective in  
said treatment.  
40

45 The disease is a disease of inflammatory, auto-immune,  
allergic, neoplastic or transplantation-related origin,  
30 or a combination of such. In accordance herewith, the  
disease may be an inflammatory bowel disease such as  
Crohn's colitis or ulcerative colitis, sclerosis, type I  
diabetes, rheumatoid arthritis, psoriasis, atopic  
50 dermatitis, asthma, malignant melanoma, renal carcinoma,  
35 lung cancer, cancer of the uterus, prostate cancer,  
hepatic carcinoma, breast cancer, cutaneous lymphoma,

rejection-related disease or Graft-versus-host-related disease.

Accordingly, candidate factors are tested in a method as described herein in place of IL-2 or IL-4 or a functionally similar compound or in addition to the combination of IL-2 and IL-4 or said functionally similar compound(s), and the effect compared to the effect obtained by using a combination of IL-2 and IL-4.

#### Methods of testing the effect of a medicament

The present invention also relates to a method of testing the effect of a medicament against a T-cell associated disease, which method comprises

- (a) providing a T-cell line as defined above,
- (b) applying the medicament to be tested to the T-cell line, and
- (c) observing the effect of the medicament on the T-cell line.

In one embodiment of this method, the cytokine profile of the T-cell line with and without the addition of the medicament is compared. Furthermore, the phenotype, proliferation and/or apoptosis of the T-cell line with and without the addition of the medicament may be compared. In particular, the intracellular amount of NF $\kappa$ B and/or JAK/STAT pathway may be monitored.

In this method, the medicament to be tested is preferably selected from compound libraries such as small molecule libraries or peptide libraries or antibodies against T-cell components. In particular, the medicament may be selected from peptide fragments from T-cell receptors.

5

Model systems

10

Thus, in a further aspect, the present invention relates to a model system for testing the effect of a medicament against a T-cell associated disease, which model system comprises at least one T-cell line as defined above.

15

Methods of detecting T-cell growth factors

20

The invention also relates to a method of detecting T-cell growth factors for use in the method of expanding and selecting disease associated T-cells as defined above, in which method candidate factors are used in place of IL-2 or IL-4 or in addition to the combination of IL-2 and IL-4, and in which the effect compared to the effect obtained by using a combination of IL-2 and IL-4.

25

15

30

Methods of monitoring responses

35

The present invention also includes a method of monitoring the response to a treatment of a disease of inflammatory, auto-immune or neoplastic origin, or combinations thereof, said method comprising comparing the phenotype, proliferation, apoptosis, and/or cytokine profile of activated T-cells in tissue sample taken from the patient to be treated before the start of the treatment and during the treatment and/or after the treatment has ended. Accordingly, this method may be used to identify patients which do not responding to a certain treatment.

45

30

Methods of identifying disease associated antigens

50

A part of the present invention is also a method of identifying disease associated antigens, comprising screening peptide libraries or antigen samples for their

55

re-activation properties in a T-cell line as defined and claimed herein.

The present invention is further illustrated by the following non-limiting examples.

#### EXAMPLES

##### EXAMPLE 1

##### Derivation of finite and continuous peripheral blood T-cell lines

Peripheral blood mononuclear cells (PBMC) from 3 healthy donors were isolated by standard Ficoll-Isopaque gradient centrifugation. The PBMC were resuspended at  $5 \times 10^5$  cells/ml in 90% RPMI 1640, 10% human AB serum, 1000u/ml IL-2 and 500u/ml IL-4 with antibiotics as described (ref. 12). To assess whether longevity of cultured PBMC is dependent on in vitro activation, PBMC were cultured in the above medium alone or with additional alloactivation.  $5 \times 10^6$  PBMC were stimulated with the heavily  $\gamma$ -irradiated (60 Gy) Psor-2 cell line at a 5:1 ratio. The Psor-2 cell line is a continuous T-cell line established from a skin biopsy specimen of a patient with psoriasis vulgaris by culturing the skin specimen in the medium mentioned above (ref. 0).

Estimation of CD28 expression as a function of cell population doublings. Monoclonal antibodies against CD3, CD4, CD8, CD28, and CD56 were purchased from PharMingen. An  $\alpha/\beta$  T-cell receptor subfamily antibody against V $\beta$ 18 was obtained from Immunotech. An indirect immunofluorescence technique was applied to label the cells as previously described (ref. 12). Allostimulated continuously growing peripheral blood T-cell lines were

cryopreserved for each 10 PD. Cells cryopreserved at different PD were then thawed, cultured for 4 days and analysed for CD28 expression by flow cytometry. CD4 and CD8 expression served as positive and negative controls, respectively. For each antibody,  $2 \times 10^4$  cells were analysed (FACS Calibur, Becton Dickinson). Fluorescence microscopy was also applied to evaluate the stainings.

A clonal CD4+, V $\beta$ 18+ T-cell line My-La, 46,XY,i(18q) (refs. 17, 18) cultured with 1000 u/ml IL-2 and 500 u/ml IL-4 was also analysed for CD28 expression at different PD.

Other methods. Cells were found to be free of mycoplasma by the Hoechst staining test. Telomerase activity of  $10^3$  cells was determined by the TRAPeze Telomerase Detection Kit as described by the manufacturer (Oncor).

Growth of peripheral blood cells with and without allostimulation. PBMC from the 3 healthy donors proliferated between 1 to 3 PD when cultured in the cytokine supplemented medium alone (Fig. 4) in agreement with previously published data showing that peripheral blood cells proliferate only transiently when stimulated with a combination of IL-2 and IL-4 (refs. 4, 12). However, when PBMC were allostimulated once with the Psor-2 cell line in the presence of a high concentration of IL-2 and IL-4, T-cells as well as non-T-cells (preferentially CD3-, CD56+) proliferated vigorously during the first 4 to 6 weeks.

After approximately 50 PD only CD4+ T-cell grew in the cytokine based medium. All three CD4+ allostimulated T-cell lines have proliferated beyond 150 PD with a PD-time of 30 to 36 hours (Fig. 4). This corresponds to an increase in cell numbers of  $2^{150} \sim 10^{45}$ -fold. As

allostimulated peripheral blood T-lymphocytes have been estimated to have a limited in vitro life-span of  $23 \pm 7$  PD (ref. 19) the allostimulated CD4+ cell lines reported here can be considered continuous, effectively having an unlimited replication capacity.

So far, the three continuous peripheral blood derived CD4+ cell lines show no sign of growth exhaustion and at PD 150 still retain alloreactivity (results not shown).

Cytokine dependent continuous T-cell lines have cytokine dependent telomerase activity. Continuous cell lines are expected to have telomerase activity. When cultured in the presence of both IL-2 and IL-4 in vitro activated peripheral blood CD4+ T-cells show high telomerase activity (Fig. 5) comparable to that of a leukemic cell line Se-Ax (ref. 20), established from a patient with Sezary's syndrome. Withdrawal of either IL-2 or IL-4 results in growth arrest. After withdrawal of IL-4, a 100 PD cell culture cease proliferating after 14 to 21 days. Withdrawal of IL-2 results in cell growth arrest between 6 to 9 days. As shown in Fig. 5 telomerase activity in IL-2 or IL-4 starved cells is severely reduced. The results indicate that simultaneous presence of IL-2 and IL-4 regulates both growth and telomerase activity in these T-cell lines.

CD28 expression correlates inversely with cell population doublings. Allostimulated PBMC cultured in the cytokine supplemented medium became pure CD4+ cell lines after approximately 50 to 60 PD. CD28 expression of one such CD4+ cell line, Act-1, at PD 60 and PD 150 is presented in Fig. 6. CD28 expression is clearly detectable at PD 60 but absent at PD 150. A gradual decline in expression of CD28 between PD 60 and PD 150 could be observed.

To investigate whether the culture system preferentially expands pre-existing CD28 negative CD4+ cells or whether CD28 could serve as a mitotic clock in individual T-cells a clonal CD4+, V $\beta$ 18+ T-cell line established from an inflammatory skin biopsy specimen (refs. 17, 18) was investigated for CD28 expression. As shown in Fig. 7, CD28 expression of this T-cell clone (My-La, 46,XY,i(18q)) decreases gradually with cell population doublings being present at PD 40 and completely absent at PD 200. However, CD4+ expression is compatible at PD 40 and PD 200. These findings are in agreement with data obtained from finite CD4+ T-cell lines (ref. 5) showing down-regulation of CD28, but not complete loss of CD28 expression with increasing PD. The results presented here show that CD28 expression correlates inversely with cell population doublings and indicates that CD28 expression can serve as a mitotic clock at the clonal level.

The results show that alloactivation with the continuous psoriatic T-cell line Psor-2 can efficiently prime allogeneric CD4+ peripheral blood T-cells to cytokine dependent continuous growth. These cytokine-driven peripheral blood derived CD4+ T-cell lines show IL-2 and IL-4 dependent telomerase activity, and they gradually lose CD28 expression with increasing cell population doublings.

Conclusion. Contrary to other normal human somatic cells T-lymphocytes can in vitro like in vivo be activated to continuous cytokine driven growth. The results presented here raises the possibility of generating an unlimited number of T-cells with predefined specificity. Such immortal T-cell lines may be useful for several applications, for instance for standardisation of T-cell mediated biological assays and for generating sufficient

5 numbers of auto-immune T-cells for human T-cell  
10 vaccination.

10 EXAMPLE 2

5  
15 Super-antigen directly augment the cytokine production of  
two novel continuous Gut-derived T-cell lines from  
patients with Crohn's disease

20 IFN $\gamma$  producing CD4 $^{+}$  T-lymphocytes have been implicated  
with progression of Crohn's disease whereas IL-10-  
producing CD4 $^{+}$  T-lymphocytes are thought to down-regulate  
disease activity.

25 15 In the following, it is investigated whether a newly  
devised cell culture protocol could select for continuous  
clonal CD4 $^{+}$  T-cell lines producing either IFN $\gamma$  or IL-10.

30 Biopsy specimens. At least eight colonic biopsies were  
20 obtained from affected mucosa of two patients. The  
biopsies were examined for histopathological changes and  
a diagnosis of Crohn's disease was established according  
35 to clinical, radiological and histopathological data.

25 In each patient, two additional biopsies were taken for  
in vitro culture of T-cells. The Gut<sub>-1</sub> T-cell clone was  
40 established from a patient undergoing cyclosporine  
treatment with a CDAI index of 296 whereas the patient  
from whom Gut<sub>-2</sub> derived had a CDAI index of 155. The  
30 study was approved by the local ethic committee.

45 Cell culture. The two biopsies were washed twice in  
sterile PBS and once in the growth medium. The growth  
50 medium consisted of 90% RPMI 1640 10% human AB serum. 100  
35 U/ml penicillin G 100 pg/ml streptomycin (basal medium,  
BM) supplemented with 2000 u/ml IL-2 and 500 u/ml IL-4



(complete medium). The T-lymphocytes were initially expanded in 5 ml complete medium and when cell density reached  $1.5 \times 10^6$ /ml, the culture was split at a 1:2 ratio.

T-cells of the primary cultures from which Gut $\alpha$ -2 derived were allostimulated with the heavily  $\gamma$ -irradiated (60Gy) leukemic cell line Se-Ax at a 5:1 ratio. The continuous Se-Ax cell line was established from a patient with Sczary's syndrome (ref. 20).

#### Phenotyping

Phenotyping. Monoclonal antibodies against CD3 (OKT3), CD4 (OKT4), CD8 (OKT8) and CD25 were obtained from hybridomas from American Type Culture Collection (ATCC). Monoclonal antibodies against CD45RO and HLA-DR were purchased from PharMingen. Monoclonal antibodies against TCR-1 (TCR $\gamma/\delta$ ), TCR-2 (TCR $\alpha/\beta$ ) and  $\alpha/\beta$  T-cell receptor subfamily antibodies against V $\beta$  1, V $\beta$  2, V $\beta$  3, V $\beta$  5.1, V $\beta$  5.2, V $\beta$  5.3, V $\beta$  7, V $\beta$  8, V $\beta$  9, V $\beta$  11, V $\beta$  12, V $\beta$  13.1, V $\beta$  13.6, V $\beta$  14, V $\beta$  16, V $\beta$  17, V $\beta$  18, V $\beta$  19, V $\beta$  20, V $\beta$  21.3, V $\beta$  22 and V $\beta$  23 were obtained through Coulter. An indirect immunofluorescence technique was applied to label the cells as previously described (ref. 12).  $2 \times 10^4$  events were analysed by flow cytometry (FACS Calibur, Becton Dickinson) and debris and aggregates were excluded by gating. Fluorescence microscopy was also applied to evaluate the stainings.

Stimulation of cells. Cells cultured in complete medium were washed twice with RPMI 1640 in order to eliminate residual cytokines. They were then re-suspended in basal medium with IL-2 or complete medium at  $10^6$ /ml. Cells were then stimulated either with 10  $\mu$ g/ml monoclonal antibodies against CD3 or with staphylococcus

enterotoxins A, B, D and E at a concentrations of 1 µg/ml (obtained from Toxin Technology Madison, WI).

Cytokine determination. Supernatant of stimulated cells and cells cultured in basal medium with IL-2 or complete medium was harvested after 24 or 48 hours. Cytokine matched antibody pairs for determination of IFN $\gamma$  IL-4, IL-10 and tumour necrosis factor (TNF $\alpha$ ) were obtained from Endogen. The detecting antibodies were all biotinylated. A time resolved fluorometric assay applying Europium labelled streptavidin and a Delphia 1234 fluorometer was used to determine the cytokine contents as described by the manufacturer (Wallac). As the cell culture medium contained human serum cytokine, concentrations below 100 pg/ml were not considered to be associated with cytokine producing T-cells. The data were analysed by a computer programme (Biosoft, Assay Zap).

Other methods. Cells were found to be free of mycoplasma by the Hoechst staining test. Karyotyping with Q banding followed standard procedures. The karyotypes were established according to the International System for Human Cytogenetic Nomenclature (ISCN) (1985).

Establishment, phenotype and constitutive cytokine production of Gut $\alpha$ -2. When placed in the complete medium, lymphocytes migrated from the biopsy specimens and proliferation was evident within a week. After approximately two weeks the cell culture had expanded to more than  $50 \times 10^6$  cells. The phenotype of this culture is shown in Fig. 8. Both TCR-1 and TCR-2 as well as CD4+ and CD8+ T-cells that are present in situ (ref. 21) are expanded in the cell culture medium. The TCR-2 population was oligo- or polyclonal as evidenced by their reaction with several V $\beta$  subfamily antibodies. A positive staining with a V $\beta$  subfamily antibody ranged from 0.2% to 8%. The

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

activation marker CD25 is only partially expressed in the growing T-cell culture (Fig. 8) and another activation marker HLA-DR differs widely in expression among individual T-cells. At this stage, the culture was split in two, half of the cells were cultured with additional allostimulation, the other half was cultured in the complete medium alone. Cells kept in complete medium without allostimulation developed into a finite cell culture dominated by CD8+ T-cells. The allostimulated culture initially also increased the percentage of CD8+ cells. However, after a period with no apparent T-cell number increase, CD4+ T-cells started to proliferate continuously. This CD4+ T-cell line Gut<sub>R</sub>-2 has proliferated beyond 250 cell population doublings (PD) with a PD time of approximately 36 hours. As allostimulated T-cell lines have been reported to have a finite life-span of 23±7 PDs, Gut<sub>R</sub>-2 can be considered immortal effectively having an unlimited replicative capacity. At PD~150 Gut<sub>R</sub>-2 became independent of IL-4 for continued growth. The phenotype of the continuous Gut<sub>R</sub>-2 cell line is presented in Fig. 9. Among the V<sub>β</sub> subfamily antibodies tested Gut<sub>R</sub>-2 only expresses the V<sub>β</sub>19 subfamily of the TCR-2 complex indicating that Gut<sub>R</sub>-2 is a clone. This assumption was confirmed by karyotyping as Gut<sub>R</sub>-2 after approximately 125 PD developed a clonal chromosome aberration observed in all metaphases (Fig. 10). Thus, also by cytogenetic criteria the V<sub>β</sub>19+ Gut-2 cell line is a clonal T-cell line. Comparison of Fig. 8 and Fig. 9 shows that clonal Gut<sub>R</sub>-2 CD4+ T-cell line develops from V<sub>β</sub>19+ T-cells that comprise less than 2% of the T-cells in the primary culture. As shown in Table 1 the V<sub>β</sub>19+ clonal Gut<sub>R</sub>-2 T-cell line constitutively produces IL-10 in basal medium with IL-2 (and also in complete medium), but without additional stimulation. IL-10 concentrations have been measured over a time period

of four months corresponding to an increase in cell numbers of approximately  $2^{80} \sim 10^{24}$  -fold.

Establishment, phenotype and karyotype of Gut<sub>1</sub>-1. Within ten days lymphocytes from the gut biopsy specimens from which Gut<sub>1</sub>-1 derived had expanded to more than  $50 \times 10^6$  cells with a phenotype distribution similar to that shown in Fig. 8. Upon culture in the cytokine based medium, but without antigen and accessory cells added, CD4<sup>+</sup> T-cells continued to expand, and within 20 PD a pure CD4<sup>+</sup> T-cell line evolved that have proliferated beyond 300 PD with a PD time of approximately 30 hours. Thus, this cell line Gut<sub>1</sub>-1 can be considered continuous. The phenotype of Gut<sub>1</sub>-1 at PD 150 is presented in Fig. 11 and, as shown, it has markers compatible with mature memory CD4<sup>+</sup> T-cells. At PD~100, Gut<sub>1</sub>-1 developed a clonal chromosome aberration as shown in Fig. 12 and like Gut<sub>R</sub>-2, Gut<sub>1</sub>-1 is also a continuous clonal CD4<sup>+</sup> cell line. By phenotyping non of the available subfamily V <sub>$\beta$</sub> , specific antibodies reacted with Gut<sub>1</sub>-1. Unlike Gut<sub>R</sub>-2 constitutive cytokine production was not detectable in Gut<sub>1</sub>-1 cells.

Super-antigens directly induce cytokine production in Gut<sub>1</sub>-1 cells and augment cytokine production in Gut<sub>R</sub>-2 cells. As Gut<sub>1</sub>-1 (and Gut<sub>R</sub>-2) expresses major histocompatibility complex class II (MHC class II) antigens that are high affinity receptors for several super-antigens, it was investigated whether these cell lines could somehow auto-present super-antigens. Four arbitrarily chosen super-antigens SEA, SEB, SED and SEE were tested for their ability to induce cytokine production in Gut<sub>1</sub>-1 cells (Table 2). As shown, soluble antibody against CD3 (OKT3) in the presence of IL-2 and IL-4 could not induce detectable cytokine production whereas SEA, SED and SEE induced IFN $\gamma$  production.

Similarly, the four super-antigens were tested for their ability to alter the cytokine production of Gut<sub>R</sub>-2 cells. As shown in Table 3, SEB induced high levels of IFN- $\gamma$  production and also significantly augmented IL-10 production in Gut<sub>R</sub>-2 cells. As SEB activation is selectively induced in T-cells bearing V $\beta$ 3,12,14,15,19 and 20 (ref. 22) the results presented in Table 3 indicate that Gut<sub>R</sub>-2 auto-present SEB as classical antigen presenting cells.

Discussion. It has been suggested that the normal tolerance to commensal intestinal bacterial antigens or super-antigens is broken in Crohn's disease. Activated CD4+ T-lymphocytes secreting IFN $\gamma$ , thereby activating monocytes/macrophages to enhanced TNF $\alpha$  production has been implicated in maintenance of Crohn's disease (ref. 23).

Gut<sub>I</sub>-1 is an inflammatory CD4+ T-cell clone established from a gut biopsy specimen without addition of mitogen, antigen and accessory cells. It is thus very likely that Gut<sub>I</sub>-1 was activated in vivo to cytokine driven growth in vitro. This assumption is compatible with the notion that inflammatory T-cells are highly activated in Crohn's disease. It should be noted that the cell culture system selects for the fastest growing T-cell clone implicating that several T-cell clones with properties like Gut<sub>I</sub>-1 exist in the inflamed gut mucosa. The V $\beta$  subfamily specificity of Gut<sub>I</sub>-1 could not be determined by phenotyping excluding the possibility of pre-selecting a super-antigen that could optimally induce IFN $\gamma$  production. However, Gut<sub>I</sub>-1 responded by direct addition of SEA, SED and SEE with IFN $\gamma$  production indicating that Gut<sub>I</sub>-1 can auto-present super-antigens. Thus, IFN $\gamma$  production by Gut<sub>I</sub>-1 cells does not necessarily require a specific antigen presented by antigen presenting cells.

5 If this property is also reflected in vivo, no specific  
microbial agent may be essential for the inflammatory  
10 response. Furthermore, inflammatory T-cells bypassing the  
classical antigen presentation could aggravate a chronic  
5 inflammation.

15 Gut<sub>R</sub>-2 is a CD4<sup>+</sup> V<sub>β</sub>19<sup>+</sup> cell clone established by  
allostimulation of outgrowing gut T-lymphocytes. During a  
period of nine months without allostimulation (150 PD)  
10 the clonal Gut<sub>R</sub>-2 cell line has constitutively produced  
20 IL-10.

As Gut<sub>R</sub>-2 expresses both high affinity receptors for SEB  
(MHC class II), and a SEB responsive V<sub>β</sub> chain (ref. 12)  
25 direct addition of SEB to Gut<sub>R</sub>-2 results in a dramatic  
15 IL-10 and IFN<sub>γ</sub> production. The cytokine production of  
activated Gut<sub>R</sub>-2 cells thus resembles a recently  
described regulatory CD4<sup>+</sup> T-cell subset (ref. 24).

30 It is intriguing to speculate that regulatory T-cells  
20 like Gut<sub>R</sub>-2 with constitutive IL-10 production  
independent of direct antigen activation may contribute  
35 to normal gut tolerance. Gut<sub>R</sub>-2 shows as mentioned above  
some properties with a newly described regulatory IL-10  
25 producing CD4<sup>+</sup> T-lymphocyte population (ref. 24).  
However, Gut<sub>R</sub>-2 differs from this sub-population by  
40 constitutive non antigen mediated IL-10 production and by  
its continuous growth.

45 An advantage of the cell culture system described here  
for gut T-cell clones is that their continuous growth  
gives rise to an unlimited number of T-cells. Such  
immortal T-cell clones may be useful for testing  
50 biological response modifiers, and inflammatory T-cell  
35 clones like Gut<sub>R</sub>-1 could provide the basis for a T-cell  
vaccination of patients with Crohn's disease.

TABLE 1. Average cytokine production (pg/ml/10<sup>6</sup> T-cells) of five different experiments between PD 150 to PD 225 of continuous growing GUT<sub>R</sub>-2 cells.

IL-4	1FN $\gamma$	IL-10	TNF $\alpha$
<100	258 (147-369)	2460 (1887-3033)	<100

Cells in basal medium with IL-2. 95% confidence intervals in parenthesis.

TABLE 2. Cytokine production (pg/ml/10<sup>6</sup> T-cells) in GUT<sub>1</sub>-1 after stimulation with superantigens (at PD 120).

GUT <sub>1</sub> -1	TNF $\alpha$	INF $\gamma$	IL-10
Complete medium	<100	<100	<100
+ antibody against CD3	<100	<100	<100
+ SEA	<100	1990 (1917-2163)	<100
+ SEB	<100	290 (164-416)	<100
+ SED	<100	1500 (1432-1568)	<100
+ SEE	<100	2070 (1910-2230)	<100

95% confidence intervals in parenthesis.

TABLE 3. Cytokine production (pg/ml/10<sup>6</sup> T-cells) in GUT<sub>R</sub>-2 after stimulation with superantigens (at PD 150).

GUT <sub>R</sub> -2	TNF $\alpha$	IFN $\gamma$	IL-10
Complete medium	<100	<100	2850 (2679-3021)
+ SEA	<100	<100	2840 (2738-2942)
+ SEB	<100	>25000	>25000
+ SED	<100	470 (453-487)	5130 (3899-5361)
+ SEE	<100	<100	5080 (4613-5867)

95% confidence intervals in parenthesis.

### EXAMPLE 3

Infliximab, a chimeric TNF $\alpha$  antibody, down-regulates the INF $\gamma$  production in activated Gut T-lymphocytes in Crohn's Disease

#### Materials and methods

Patients. The biopsy specimen were obtained from 5 patients with an established diagnosis of Crohn's disease according to clinical, radiological and histopathological criteria (1 male 22 years, and 4 females, mean: 38 years, range: 34-43 years). All the patients had active disease with a CDAI index above 150.

Biopsy specimens. Two colonic biopsies were obtained from each anatomical segment of the affected mucosa in each patient during colonoscopy (in total 16 biopsies). The biopsies were evaluated for histopathological changes. In



each patient T-cells were cultured from four additional biopsies from mucosa with macroscopically active disease. The study was approved by the local ethic committee of Aarhus County.

Cell culture. The four biopsies were washed twice in sterile PBS (saline) and once in the growth medium. The growth medium consisted of 90% RPMI 1640 10% human AB serum. 100 U/ml penicillin G 100 µg/ml streptomycin (basal medium, BM) supplemented with 2000 u/ml IL-2 and 500 u/ml IL-4 (complete medium). The T-lymphocytes were initially expanded in 5 ml complete medium and when cell density reached  $1.5 \times 10^6$ /ml, the culture was split at a 1:2 ratio. From one female, two cultures were established from specimen taken 8 month apart (C1x and C11.3), and from the male two cultures were established from two different anatomical lesions (one from the cecum and one from the descending colon (C8.1 and C8.3 respectively). In the remaining three patients, one representative culture was used for the experiments. C1x, C2x and C4.2 are cultures grown for more than 150 days without further addition of antigen or feeder cells. C11.3, C12.1, C8.1, C8.3 are primary cultures cultured for less than 50 days.

Phenotyping and transmembrane TNFα. Monoclonal antibodies against CD3 (OKT3), CD4 (OKT4), CD8 (OKT8) and CD25 were obtained from hybridomas from American Type Culture Collection (ATCC). Monoclonal antibodies against CD45RO and HLA-DR were purchased from PharMingen. Monoclonal antibodies against TCR-1 (TCRα,β), TCR-2 (TCRγ,δ) and T cell receptor subfamily antibodies against V<sub>β</sub>-chains were obtained through Coulter. An indirect immunofluorescence technique was applied to label the cells as previously described (ref. 12).  $2 \times 10^4$  events were analysed by flow cytometry (FACS Calibur, Becton Dickinson) and debris and aggregates were excluded by gating. Fluorescence

microscopy was also applied to evaluate the staining. The antibody used for detection of transmembrane TNF $\alpha$  was obtained from R&D (FAB210 FITC).  $5 \times 10^5$  cells were obtained. 15  $\mu$ l of undiluted antibody was added for 45 minutes. Unbound antibody was removed by washing,  $2 \times 10^5$  cells were analysed by flow cytometry (FACS). The binding of Infliximab was determined in an indirect way by a competitive assay with untreated cells as control.

In vivo activated primary cultures. These cells were washed once in RPMI 1640. They were then re-suspended in complete medium with and without Infliximab. Infliximab (obtained from Centocor, Malvern, Pennsylvania) was added in a concentration of 5  $\mu$ g/ml cell culture. Transmembrane TNF $\alpha$  and apoptosis was detected after one hour and 24 hours.

SEA stimulation of primary culture (C8.3) and cultures grown for more than 150 days (C1x, C2x, C4.2). Cells used had been cultured in complete medium. The cells were washed twice with RPMI 1640 in order to eliminate residual cytokines. They were then re-suspended in complete medium at a cell density of  $10^5$ /ml. Cells were then stimulated with Staphylococcus enterotoxins A (SEA) (obtained from Toxin Technology Madison, WI) at a concentration of 0.5  $\mu$ g/ml cell culture. Two hours after stimulation cells were washed twice in RPMI 1640 and then re-suspended in complete medium with or without Infliximab (at a concentration as described previously). Transmembrane TNF $\alpha$  was determined one hour and 24 hours after the addition of Infliximab in activated cells and controls.

Cytokine determination. Supernatant of stimulated cells and controls was harvested after 24 hours. Cytokine matched antibody pairs for determination of IFN $\gamma$  and

5 tumour necrosis factor (TNF $\alpha$ ) were obtained from Endogen.  
The detecting antibodies were all biotinylated. A time  
10 resolved fluorometric assay applying Europium labelled  
streptavidin and a Delphia 1234 fluorometer was used to  
5 determine the cytokine contents as described by the  
manufacturer (Wallac). Briefly, plates were covered with  
15 50  $\mu$ l of coating antibody at a concentration 2.5  $\mu$ g/ml.  
They were placed at 4°C overnight. Afterwards they were  
blocked with 10% AB-serum. Supernatant, controls and  
20 standards were added for two hours. Biotinylated antibody  
was added at a concentration of 1  $\mu$ g/ml for one hour.  
Addition of Eu<sup>3+</sup> marked streptavidin at a concentration  
1:2000. Addition of enhancement solution. After 20  
25 minutes, the plates could be read at a Delfia  
fluorometer. Because cytokine instability in low  
concentrations, new standards and dilutions were  
established for each determination. As the cell culture  
medium contained human serum, medium was controlled for  
30 cytokine content and levels were used as background.  
Concentrations below 30 pg/ml were not considered to be  
associated with cytokine producing T-cells. The data were  
analysed by a computer program (Assay Zap, Biosoft).  
35 Values were averages of three determinations.

25 Apoptosis and cytolysis. Annexin-FITC and propidium  
iodide were used for the determination of apoptosis  
40 (Nexins research and R&D).  $5 \times 10^5$  cells were obtained and  
placed in buffer for a half hour. Half of the cells were  
stained with 5  $\mu$ l of Annexin-FITC diluted 1:10 in buffer,  
30 and 2.5  $\mu$ l of propidium iodide. After incubation for 15  
minutes the cells were analysed on a flow cytometer (FACS  
Calibur, Becton Dickinson). For the determination of  
cytolysis the same procedure was used after the addition  
50 of Infliximab and incubation with fresh human serum for  
35 one hour. A murine HLA class II antibody was used as a  
positive control for Infliximab.

Proliferation. Cell cultures were monitored with a Coulter counter measuring the increment in cell count after 24 hours. The channelysed count is measured on a 500 µl test sample. It was diluted 40 times in 20 ml Isoton II<sup>®</sup> (Coulter), so the cell count/ml was 80 times the channelysed count/ml.

Other methods. Cells were found to be free of mycoplasma by the Hoechst staining test.

### Results

Cell culture and phenotype. When placed in the complete medium, lymphocytes migrated from the biopsy specimens and proliferation was evident within a week. After approximately two weeks the cell culture had expanded to more than  $50 \times 10^6$  cells. No antigen nor feeder cells were added. The in vivo activated T-cells were expanded only in the presence of high concentrations of IL-2 and IL-4. The phenotype of the primary cultures (C11.3, C12.1, C8.1, C8.3) is shown in Fig. 13 (representative example). Both TCR-1 and TCR-2 as well as CD4<sup>+</sup> and CD8<sup>+</sup> T cells that are present in situ are expanded in the cell culture medium. Upon continued culture a pure CD4<sup>+</sup> cell line evolved within 40-50 days. C1x, C2x and C4.2 are representatives that have proliferated beyond 150 days. The cultures described above were used to study the effects of Infliximab on cytokine production, transmembrane TNF $\alpha$ , apoptosis, cytolysis and growth.

Cytokine production. In all the primary cultures a spontaneous production of IFN $\gamma$  was observed. In all cultures, Infliximab induced a reduction in the 24 hour production of IFN $\gamma$  (Fig. 14). As a control, recombinant IFN $\gamma$  was added to the supernatant together with

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

Infliximab (0.5 ng/ml recombinant IFN $\gamma$  and 5  $\mu$ g/ml Infliximab). The triple determination of IFN $\gamma$  was 0.45 ng/ml. The TNF $\alpha$  productions in the primary cultures were close to detection level (<50 pg/ml), but this production was markedly enhanced by the stimulation with super-antigen (Fig. 15).

15  
20  
25  
30  
35  
40  
45  
50  
55

In the SEA stimulated primary culture, the effect of Infliximab on the absolute cytokine production was more pronounced (C8.1, 26 days, IFN $\gamma$ : 25 to 12.81 ng/ml (49%), TNF $\alpha$ : 1.95 to 0.05 (97%); C8.3, 35 days, IFN $\gamma$ : 10.85 to 4.78 (56%) TNF $\alpha$ : 11.9 to 0.3 (97%)) the reduction without stimulation with SEA was C8.1, IFN $\gamma$ : 0.16 to 0.05 (68%), (TNF $\alpha$  below detection limit); C8.3: IFN $\gamma$ : 2.58 to 1.47 (43%), (TNF $\alpha$  below detection limit).

30  
35  
40  
45  
50  
55

In the cultures grown for more than 150 days, there was not any constitutive production of IFN $\gamma$ , but after stimulation with SEA an increase in the production of IFN $\gamma$  and TNF $\alpha$  was observed. This cytokine production was also reduced by the addition of Infliximab (Fig. 16). There was no correlation between the level of IFN $\gamma$  or TNF $\alpha$  production and the amount transmembrane TNF $\alpha$ .

25  
30  
35  
40  
45  
50  
55

Membrane bound TNF $\alpha$  and binding of Infliximab. The primary cultures all presents transmembrane TNF $\alpha$  determined by FACS analysis (Fig. 13). After addition of Infliximab to the cultures, the staining intensity of transmembrane TNF $\gamma$  is reduced indicated by a left shift of the FACS curve. If these primary cultures were stimulated by super-antigen (SEA), an increase in the amount of transmembrane TNF $\alpha$  was observed, and the difference after supplement of Infliximab was more evident.

35

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

In the cell lines C2x, C1x and C4.2 transmembrane TNF $\gamma$  was evident after stimulation with super-antigen SEA and Infliximab affected this relationship. No activation (indicated by transmembrane TNF $\alpha$ ) could be demonstrated in these long term grown cultures before the addition of SEA (Fig. 17).

Apoptosis and cytolysis. FACS analysis of Annexin-FITC and propidium iodide stained cells was used as a measure of apoptosis and necrosis with and without the addition of complement. As a positive control a HLA-class II antibody was used.

Infliximab did not induce any apoptosis in any of the in vivo activated primary cultures. In the long term cultured SEA stimulated C2x, Infliximab did not increase neither the amount of propidium iodide nor Annexin-FITC positive cells compared with SEA alone (Fig. 18) (HLA class II antibody as control).

Proliferation. Proliferation was measured by a Coulter particle counter. Infliximab did not change the proliferation rate in any of the cultures (Fig. 19 A, B and C) (primary culture)). Cultures activated by SEA gave identical results.

Discussion. T-cell activation in Crohn's disease is one of the cornerstones in the inflammatory process with epithelial destruction (refs. 25, 26), probably because the production of pro-inflammatory Th1 cytokines INF $\gamma$  and TNF $\alpha$  is increased (ref. 27).

Recent clinical studies in patients with Crohn's disease have demonstrated dramatic clinical responses following treatment with chimeric TNF $\alpha$  antibody (Infliximab) (refs. 7, 9). Different mechanisms have been proposed. Decreased

5  
10 production of TNF $\alpha$  by T-cells and the neutralisation of circulating TNF $\alpha$  may indirectly reduce the production of IFN $\gamma$  (ref. 28).

15 5 Animal studies have demonstrated that transmembrane TNF $\alpha$  in the genetically engineered SP2/O myeloma cell line can bind Infliximab activating complement and macrophages resulting in cytolysis. In the present study, we have described the in vitro effects of Infliximab on in vivo  
20 10 activated T-cells obtained from the colon of patients with active Crohn's disease, regarding production of IFN $\gamma$  and TNF $\alpha$ , binding to transmembrane TNF $\alpha$ , apoptosis and proliferation. Infliximab down-regulates the IFN $\gamma$  and TNF $\alpha$  production in all primary T-cell lines. These  
25 15 cultures revealed a spontaneous production of IFN $\gamma$  and to a lesser extent TNF $\alpha$ . This type 1 cytokine profile indicates that the primary cultures are in vivo activated since no antigen nor feeder cells has been added in  
30 20 vitro. In a previous study, it has been demonstrated that TNF $\alpha$  may be necessary for the LPMC production of IFN $\gamma$  (ref. 28). Although not all the primary T-cell lines did  
35 25 produce detectable amounts of TNF $\alpha$ , Infliximab reduced the IFN $\gamma$  production, probably by other mechanisms not involving TNF $\alpha$  synthesis. In cultures grown for more than  
40 30 150 days, no residual in vivo derived antigen stimulation was present illustrated by the fact, that these cultures did not have any constitutive cytokine production. After  
45 35 SEA stimulation, a pro-inflammatory cytokine profile was present illustrated by increase in the production of IFN $\gamma$  and TNF $\alpha$ . Infliximab reduced the synthesis of both IFN $\gamma$  and TNF $\alpha$ . No correlation was observed between the level  
50 40 of reduction in IFN $\gamma$  and TNF $\alpha$ . Spontaneous or stimulated secretion of IFN $\gamma$  and TNF $\alpha$  in T-cells isolated from the mucosa of patients with Crohn's disease has been closely  
55 45 related to the degree of inflammation (refs. 9, 26, 28, 29), and the levels of TNF $\alpha$  secretion in pokeweed mitogen

5 stimulated early cultures from patients with Crohn's  
disease in remission has also been related to the risk of  
10 relapse (ref. 30). The present in vitro data supports the  
clinical data. A decrease in disease activity in  
5 Infliximab treated patients would be expected if the INF $\gamma$   
and TNF $\alpha$  production is reduced in the activated  
15 intestinal T-cells.

Transmembrane TNF $\alpha$  is present in the primary cultures.  
10 The presence of transmembrane TNF $\alpha$  indicates a state of  
in vivo T-cell activation as illustrated previously (ref.  
20 31). It has been shown that transmembrane TNF $\alpha$  correlates  
with the expression of the activation marker CD69. This  
finding is in good agreement with the Th1 cytokine  
25 profile in these cell lines. Only a minor fraction of the  
T-cells in the cultures are activated, but after addition  
of Infliximab, a left shift in the FACS curve is observed  
indicating the binding of Infliximab to the T-cells. If  
30 the culture is stimulated by super-antigen, the  
activation is more pronounced and the binding of  
20 Infliximab is demonstrated more clearly. In cultures  
grown for more than 150 days, no residual in vivo derived  
35 antigens are present. In these T-cell lines, no  
transmembrane TNF $\alpha$  could be demonstrated. After SEA  
25 stimulation, transmembrane TNF $\alpha$  was prominent, and a  
Infliximab-induced competitive inhibition could be shown.  
40 The 26KD transmembrane TNF $\alpha$  is a co-stimulatory factor in  
the activation of B-cells (ref. 31). Inhibition of  
costimulatory signals by Infliximab binding to  
45 30 transmembrane TNF $\alpha$  may be of importance, and since only  
activated T-cells presents transmembrane TNF $\alpha$ , this  
effect may be confined to pro-inflammatory activated T-  
cells.

50 35 The Infliximab-induced reduction in cytokine production  
may be a result of a change in intracellular T-cell



5  
10 signalling either by a direct effect of binding to  
transmembrane TNF $\alpha$  or by an indirect effect because of  
changes in co-stimulation and T-cell interaction.  
15  
20  
25  
30  
35  
40  
45  
50  
55  
Infliximab probably binds to other epitopes of  
transmembrane TNF $\alpha$  than the FAB 210F antibody.  
Substantial evidence is the fact that Infliximab  
neutralises circulating TNF $\alpha$  which FAB210F does not so  
simple correlation between the blocking effects can not  
be established.

10  
20 In murine SP2/O myeloma cells, complement could be  
activated by binding Infliximab. This type of  
transmembrane TNF $\alpha$  was different from the wild-type by  
lacking two amino acids and a Ala in place instead of  
25 Val. This transmembrane TNF $\alpha$  was resistant to proteolytic  
cleavage. Scatchard analyses showed the cells of interest  
bound about 35000 Infliximab molecules per cell. We could  
not confirm these results in human T-cell lines. This  
30 might be related to proteolytic cleavage of human  
transmembrane TNF $\alpha$  or less extensive binding of  
20 Infliximab to in vivo activated human T-cells.

35  
40 Apoptosis may be induced in response to various cytotoxic  
stimuli including activation of cell surface receptors  
25 such as Fas or TNFR1. The ligand for the transmembrane  
TNF $\alpha$  is not fully understood, but substantial evidence  
supports the hypothesis that the co-stimulatory signals  
are mediated by the p55 subunit (TNFR1) and not the p75  
subunit (TNFR2) (ref. 31). We could not demonstrate any  
45 increased or decreased apoptosis by the binding of  
30 Infliximab to the transmembrane TNF $\alpha$  in any cultures.

50 Proliferation was unaffected in the cell lines ligated  
with Infliximab compared to the untreated cell lines. In  
35 clinical studies (ref. 28), patients responding to

5 treatment with Infliximab disclosed reduced numbers of  
10 LPMC after a single dose.

10 In summary, we found that when activated T-cells binds  
5 Infliximab the production of the pro-inflammatory  
cytokines IFN $\gamma$  and TNF $\alpha$  is reduced. Infliximab binds to  
15 transmembrane TNF $\alpha$  in activated human intestinal T-cells,  
and the binding is related to the level of activation  
demonstrated by FACS analysis and cytokine assays. We  
20 could not support results in murine myeloma cell lines  
where the binding of Infliximab activates complement  
resulting in cell lysis. Apoptosis and proliferation was  
unaffected by Infliximab. Changes in co-stimulatory  
25 signals via the TNFR-I might be a possible mechanism by  
15 which Infliximab exerts its effects.

#### EXAMPLE 4

##### 30 Examples of T-cell vaccination

##### 20 A. Multiple sclerosis (MS)

35 1. A convenient amount, for example 50 ml, blood in hepa-  
rin is drawn from a patient with MS.

25 2. The mononuclear cells of the blood that, other than  
40 lymphocytes, contain antigen presenting cells (APC) are  
isolated by a standard Ficoll-Isopaque gradient hydro-  
extracting.

45 30 3. The cells are disseminated in for example five cultur-  
ing bottles in the medium consisting of 90% RPMI 1640,  
10% human AB serum, antibiotic as well as 1000 u/ml IL-2  
50 and 500 u/ml IL-4. If convenient other cytokines as GM-  
35 CSF and TNF $\alpha$  can be added to the bottles to

increase/promote the maturing of the dendritic cells with a strong antigen presenting function.

At this stage, a selection for antigen activation (i.e. CD69+) may be included.

4. On day 0 antigen, in this case myelin, that the patient's auto-reactive T-cells react against, is added to one of the bottles. Instead of myelin components of the myelin can be added such as myelin basic protein or proteolipid protein or immune dominating epitopes deriving from these proteins.

5. This addition of antigen is repeated in the next bottle for example on day 2 and the procedure is continued with the other bottles with an interval of a couple of days.

6. Subsequently, the cells are propagated in the IL-2 and IL-4 containing medium. Notice that if "only" the life of the T-lymphocytes can be increased from 23 PD to 60 PD instead of  $10^7$  cells one will have/get  $2^{60}-10^{18}$  cells, the equivalent of 1000 tons of cells, which will be sufficient to continue all further experiments and vaccination. In case the T-cells apparently does not have the expected ability for growth the antigen stimulation can be repeated, and furthermore co-stimulation with for example phorbol ester or mitogen-stimulation may be tried to increase the growth potential.

7. The T-cells are tested for their antigen specificity and will after activating and attenuating (e.g. by  $\gamma$ -radiation, 60 Gy) be ready for T-cell vaccination.

8. Vaccination can be accomplished with  $100-500 \times 10^5$  T-cells in each forearm subcutaneously.

5

B. Insulin dependent diabetes

10

The same procedure as for A can be used, if only the  
5 antigen is for example glutamin acid decarboxylase (GAD)-  
65, GAD-67, insulin, or heat shock protein 60 (Hsp60).

15

C. Crohn's disease and ulcerative colitis

20

10 Crohn's disease is a multifactorially conditioned chronic  
inflammatory intestinal disease where the normal  
tolerance of the immune system to the microbial  
intestinal flora is broken. Here the immune reactive T-  
cell clones (for T-cell vaccination) against the  
25 microbial flora can be brought about in the following  
way:

30

From a intestinal biopsy the aerob as well as the anaerob  
bacteria are cultured. After the culturing they are  
20 sonicated and can now be used as antigen/super-antigen.  
Subsequently, the biopsy is washed in a antibiotic-  
containing medium, and within 14 days the T-lymphocytes  
35 from the biopsy can be propagated in large number  
( $>50 \times 10^6$ ) in an IL-2 and IL-4-containing medium. Antigen  
25 presenting cells are obtained by ficoll separation of the  
patient's blood cells, and antigen specific/super-antigen  
40 specific continuous intestinal T-cell clones can now be  
propagated by adding antigen and  $\gamma$ -irradiated antigen  
presenting cells to the intestinal biopsy T-cells.

45

30 An analogous strategy can be used for patients with  
ulcerative colitis.

50

Note that for procedure A and B as well as for procedure  
35 C, the vaccination is individual (depending on the type  
of tissue), i.e. it has to be the patient's own cells

55

that are used. Besides, note that T-cell vaccination primarily has been intended for persons that are already affected by diseases.

Activation (7 above) may be accomplished by mixing with a sonicated faeces sample from the patient. Such sample will contain the antigen that initially activated the T-cells in vivo. Therefore, the sonicate is suitable for boosting the T-cell lines prior to administration.

In a further alternative according to the invention in general, the biopsy or cell sample is cultured comprising IL-2 and IL-4 to enrich for activated T-cells, and the activated T-cells is isolated by immunomagnetic beads separation methods. The separated activated T-cells (which are often alloreactive) are then allostimulated and is further cultured in the presence the cytokines mentioned above. Hereby the alloactivated T-cells are expanded resulting in an antigen specific T-cell line.

This procedure may be used for any other relevant disease including the diseases mentioned above.

#### EXAMPLE 5

Establishment and characterisation of in vivo activated T cell lines from patients with Crohn's disease in preparation for immune therapy

Aim. T-cell vaccination (immunisation with attenuated auto-reactive T-cells) could be an attractive treatment option in patients with Crohn's disease. T-cell vaccination has not hitherto been possible, because auto-reactive T-cells have (like other human T-cells) limited replicative capacity in vitro (cellular senescence).

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

With the cell culture system described herein, it has been possible in certain situations, to expand and select in vivo activated T-cells in unlimited amounts. With this project we want to investigate whether such in vivo activated T-cells established from intestinal biopsies from patients with Crohn's Disease has reactivity against the patients own microflora, and if such T-cells could be used as a T-cell vaccination.

10 Background. Different studies has rendered that Crohn's  
20 disease is a multifactorial determined auto-immune  
disease where the normal tolerance against the microbial  
flora in the intestine is broken. The reactivity against  
25 the intestinal flora is mediated by reactive T-cells  
15 producing IFN $\gamma$  and TNF $\alpha$ , and these cytokines contribute  
to the destruction of the intestinal mucosa (auto-immune  
reaction) in the diseased bowel. Treatment of Crohn's  
disease has lately been concentrated on interference with  
30 the immune response by using IL-10 or TNF $\alpha$  antibodies.

20  
35  
40  
45  
50  
55

Animal experiments in murine models for auto-immune  
disease has demonstrated that immunisation with  
attenuated auto-antigen reactive T-cell clones (T-cell  
vaccination) was an effective treatment against these  
25 diseases. It has been hypothesised that the auto-reactive  
T cell clones, often with a Th-1 cytokine profile  
(producing IFN $\gamma$  and TNF $\alpha$ ) activates regulatory T-cells  
(IL-10 producing) in the immunological network.  
Regulatory T-cells are specifically directed against  
30 auto-reactive T-cells, and the production of IL-10 and  
TGF $\beta$  is immuno-suppressive to the auto-reactive cells and  
the bystander T-cells contributing in the auto-immune  
process. The advantage of T-cell vaccination to  
systemically treatment with IL-10 or TGF $\beta$  is that the  
35 regulatory T-cells are activated locally at the scene of  
inflammation and not associated with systemic adverse

5 events. Besides, it is possible that T-cell vaccination  
10 activates other effector mechanisms in the immunological  
network, as e.g. cytotoxicity, against the auto-reactive  
T-cells.

5 In the murine experiments, auto-reactive T-cell lines  
15 used for vaccination have the advantage that they are  
continuous (immortal) resulting in unlimited amounts of  
T-cells available for the relevant studies. So far, it  
20 has been postulated that human T-lymphocytes are  
restricted by cellular senescence respecting the Hayflick  
limit ( $23 \pm 7$  cell population doublings (PD)), one T-cell  
clone can expand to  $2^{23} \approx 10^7$  T-lymphocytes. This amount is  
too little for a human T-cell vaccine.

25 Preliminary results. In certain situations, T-cells do  
not respect cellular senescence in vitro. We have shown  
30 that T-lymphocytes from patients with inflammatory skin  
diseases can be cultured continuously in a medium  
20 supplied with IL-2 and IL-4 but without antigen or  
accessory cells added (ref. 4, 12, 18). These immortal T-  
cell lines are activated in vivo in a way so they can be  
35 grown in vitro with unlimited replicative capacity.  
Recently a in vitro method has been demonstrated where T-  
25 lymphocytes can be immortalised in the presence of  
antigen and IL-2 and IL-4, cf. Example 1.

40 If the replicative capacity of T-cells can be increased  
from 30 PD to 50 PD, the amount of T-cells will increase  
45 from  $2^{30} \approx 10^9$  cells (equivalent to 1g cells) to  $2^{50} \approx 10^{15}$   
cells (1 ton cells). T-cell clones are usually expanded  
by using mitogen and radiated mononuclear cells or EBV  
immortalised B-lymphoblasts. None of these methods using  
50 feeder cell populations can immortalise human T  
35 lymphocytes, cf. Example 1.

5 Recently we have demonstrated that in vivo/in situ  
activated gut T-lymphocytes from patients with active  
10 Crohn's disease and with inducible INF $\gamma$  production can be  
expanded in unlimited numbers (cf. Example 2). Such CD4+  
5 T-cells expresses besides the T-cell receptor HLA Class-  
II antigens, and can auto-present super-antigens  
15 resulting in production of large amounts of INF $\gamma$  (cf.  
Example 2). These in vivo/in situ activated CD4+ T-cells  
with a type 1 cytokine profile are probably "auto-  
10 reactive" inflammatory T-lymphocytes. Preliminary results  
shows, that the inflammatory T-cells can activate  
20 regulatory CD4+ T-cells producing IL-10 indicating that  
the established inflammatory CD4+ T-cells could be used  
for T-cell auto-vaccination in patients with Crohn's  
25 disease.

Future Studies. We have established three inflammatory  
and corresponding three regulatory autologous continuous  
30 T-cell lines from gut biopsies of patients with Crohn's  
disease. The results are the substance in a protocol  
20 which probably can be used to develop in situ activated  
T-cells in unlimited amounts from most patients with  
Crohn's disease. One of the goals in the coming studies  
35 is to expand the inflammatory T-cells according to this  
protocol in a larger number of patients.  
25

40 Activation of the inflammatory T-cells with  
antigens/super-antigens from the patients own intestinal  
flora will be of importance in the evaluation of the  
suitability of the cells as a T-cell vaccination. If they  
30 increase the production of type 1 cytokines after  
activation it indicates that the T-cells are auto-  
45 reactive. Antigen and super-antigen is obtained by  
cultures (aerobically and anaerobically) from rectal  
50 mucosa according to the methods described by Duchmann  
35 (ref. 6). As antigen presenting cells are used autologous



5 PBMC or dendritic cells. It will be studied whether  
immune-modulating (immune down-regulating) drugs inhibits  
10 the pro-inflammatory response after activation  
antigen/super-antigen. Infliximab (chimeric TNF $\alpha$   
5 antibody), 5-ASA, and steroids will be drugs of interest.

15 The interaction between autologous inflammatory and  
regulatory T-cells with and without externally activation  
will be analysed to describe if the type-/ideotype  
10 response has any implication in the activation of the  
regulatory T-cells when inflammatory T-cells are present.  
20

Perspectives. In situ/in vivo activated T-lymphocytes  
from gut biopsies of patients with Crohn's Disease has a  
25 CD4+ phenotype and a cytokine profile (IFN $\gamma$ ) that is  
compatible with a "auto-reactive" origin. If it can be  
demonstrated that these continuous T-cell lines have  
reactivity against the patients own microflora, a T-  
30 cell/T-cell receptor peptide vaccination will be a  
potential option in these patients. If such a treatment  
20 has a positive effect, perhaps curative, it could be an  
option in other auto-immune diseases as multiple  
35 sclerosis and insulin dependent diabetes mellitus and  
inflammatory diseases as psoriasis, atopic dermatitis and  
25 rheumatoid arthritis.

40 The study has been approved by the Local Ethical  
committee of Aarhus County J. nr. 1997/3855, 1997/3856,  
1998/4330, 1998/4419.

#### 30 EXAMPLE 6

##### Cancer

50 Most cancers are associated with tumour infiltrating  
35 lymphocytes (TIL), and these TIL's are known to have  
killer cell activity against the tumour cells.

One example of cancerous diseases which could be treated with the T-cell lines or T-cells prepared according to the present invention is metastatic malignant melanoma.

The procedure could be as follows:

A cutaneous biopsy specimen or a lymph node biopsy specimen is known to harbour TIL's. The biopsy is divided into two, one part being cultured without cytokines in order to establish a tumour cell line (Fig. 20A). From the other part, T-lymphocytes are expanded in a medium supplemented with e.g. 10% human AB serum, 10 nM IL-2 and 2.5 nM IL-4 in the presence of 100  $\mu$ M of the caspase inhibitor Z-VAD.

The outgrowing T-lymphocytes are in general of oligoclonal origin and consist of both CD4+ and CD8+ T-lymphocytes. Contained within the latter population are the presumed auto-immune effector cells (killer cells), while contained within the former population are CD4+ cells mediating help in generating CD8+ effector cells.

Following expansion, appropriate selection procedures may be used to select for CD8+ cells with tumour cell reactivity. Fig. 20B shows the result 24 hours after mixing an expanded CD8+ oligoclonal culture comprising cytotoxic cells with cytotoxic activity against autologous melanoma cells with melanoma cells. It should be noted that continuous T-cell lines are often oligoclonal for more than 100 PD, implying that continuous CD8+ tumour specific T-lymphocyte cell lines may react with several melanoma associated antigens, thus minimising the risk of tumour escape.

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

Selection for melanoma specific CD8+ cells may also be obtained by mixing outgrowing T-lymphocytes with tumour cells in a medium with IL-2, IL-4 and Z-VAD, because the tumour cells (target cells) acts as antigen presenting cells by directly presenting tumour associated peptides to CD8+ T-lymphocytes.

10  
20  
25  
30  
35  
40  
45  
50  
55

The tumour specific CD8+ T-lymphocytes are  $\gamma$ -irradiated in order to ensure that the cells cannot divide further and infused into the patient according to an established malignant melanoma IL-2 therapy protocol as already used by practitioners. Before administration, e.g. infusion, the T-lymphocytes can be incubated with the caspase inhibitor Z-VAD, in order to reduce AICD, or Z-VAD may be given during the administration.

30  
35  
40  
45  
50  
55

Production of these cytokines together with IFN $\gamma$  has consistently been found in 12 outgrowing T-lymphocyte cultures established from biopsies of patients with melanoma.

## REFERENCES

1. WO 88/07077 (The Children's Hospital, Inc.)
2. WO 94/02156 (The Board of Trustees of Leland Stanford Junior University)
3. WO 97/05239 (Celltherapy)
4. Kaltoft K et al., Cancer Genet. Cytogenet. 85, 68-71 (1995)
5. Effros RB, Pawelac PS, Exp. Cell Res. 25, 1129-1133 (1995)
6. Durchmann R et al., Clin. Exp. Immunol. 102, 448-455 (1995)
7. Dullemen HM et al., Gastroenterol. 109, 129-135 (1995)
8. Stronkhorst A et al., Scand. J. Gastroenterol. 27 Suppl., 61-65 (1992)
9. Targan SR et al., N. Engl. J. Med. 337, 1029-1035 (1997)
10. Groux et al.,
11. Gorrigan GJ et al., Am. J. Resp. Cell Mol. Biol. 12, 567-576 (1995)
12. Kaltoft K et al., Arch. Dermatol. Res. 287, 42-47 (1994)

13. Rolfe FG et al., Am. J. Resp. Cell Mol. Biol. 17,  
243-250 (1997)

14. Madaer R et al., Lancet 346, 807-808 (1995)

15. Zang J et al., Science 261, 1451-1454 (1993)

16. Rosenberg SA et al., JNCI 86, 1159-1166 (1994)

17. Kaltoft K et al., In Vitro Cell Dev. Biol. 28A, 161-  
167 (1992)

18. Kaltoft K et al., Cancer Genet. Cytogenet. 81, 13-16  
(1995)

19. Perillo NL et al., Exp. Geront. 24, 177-187 (1989)

20. Kaltoft K et al., Arch. Dermatol. Res. 279, 293-298  
(1997)

21. Christien J et al., Clin. Exp. Immunol. 73, 63-69  
(1987)

22. Cottrez F et al., J. Immunol. Methods 172, 85-94  
(1994)

23. Broux H, Nature 389, 737-742 (1997)

24. Pavelec et al., Immunological Review 16D, 31-42  
(1997)

25. Lionetti P et al., Gastroenterol. 105, 373-381 (1993)

26. Pender SL et al., Gut 39, 284-290 (1996)

27. Deem et al., Clin. Exp. Immunol. 83, 79-84 (1991)

5

28. Plevy SE et al., J. Immunol. 159, 6276-6282 (1997)

10

29. Schreiber SL et al., Immunol. Today 13, 136-142  
5 (1992)

15

30. Schreiber SL et al., Lancet 353, 459-461 (1999)

31. Aversa G et al., J. Exp. Med. 177, 1575-1585 (1993)

20

25

30

35

40

45

50

55

**Claims**

5

10

15

20

25

30

35

40

45

50

55

5

## CLAIMS

10

1. A method of expanding and selecting disease associated T-cells comprising

5

15

(a) obtaining a tissue sample from a mammal including a human being, the sample comprising disease activated T-cells, or

20

obtaining T-cells and antigen-presenting cell from said mammal and mixing said cells with a disease associated antigen or antigens, and

25

(b) culturing said tissue sample or said mixture of cells and antigen(s) in the presence of at least two factors which promote T-cell growth and optionally one or more additional compounds.

30

2. A method according to claim 1, wherein the factors which promote T-cell growth are selected from the group consisting of cytokines which promote T-cell growth.

35

3. A method according to claim 2, wherein the cytokines are selected from the group consisting of IL-2, IL-4, IL-7, IL-9, IL-10, IL-15, IL-16 and functionally similar cytokines.

40

45

4. A method according to any one of claims 1-3, wherein a combination of IL-2 and/or IL-15 and IL-4 and/or IL-7 and/or IL-9 is used.

50

5. A method according to any of claims 1-4, wherein a combination of IL-2 and IL-4 is used.

55

6. A method according to any one of claims 1-5, wherein each of the cytokines is used in a concentration of at



5 least 1 nM, preferably more than 2.5 nM, more preferably more than 10 nM.

10 7. A method according to any one of claims 1-6, wherein  
5 the tissue sample is selected from a biopsy, from sputum, swabs, gastric lavage, bronchial lavage, intestinal  
15 lavage, or body fluids such as spinal, pleural, pericardial, synovial, blood and bone marrow.

20 8. A method according to any one of claims 1-7, wherein  
20 the disease associated T-cells are CD4+, CD8+ or CD4-  
/CD8- T-cells.

25 9. A method according to any one of claims 1-8, wherein  
25 the disease associated T-cells are selected from the group consisting of inflammatory, cytotoxic and  
regulatory T-cells.

30 10. A method according to any one of claims 1-9, wherein  
20 the disease associated T-cells are associated with a disease of inflammatory, auto-immune, allergic,  
neoplastic or transplantation-related origin, or  
35 combinations thereof.

25 11. A method according to claim 10, wherein the disease  
of inflammatory or allergic origin is a chronic  
40 inflammatory disease, or a chronic allergic disease.

45 12. A method according to any one of claims 1-11, wherein  
30 the disease is an chronic inflammatory bowel disease, such as Crohn's disease or ulcerative colitis, sclerosis,  
type I diabetes, rheumatoid arthritis, psoriasis, atopic  
dermatitis, asthma, malignant melanoma, renal carcinoma,  
50 breast cancer, lung cancer, cancer of the uterus,  
35 prostatic cancer, cutaneous lymphoma, hepatic carcinoma,

rejection-related disease, or Graft-versus-host-related disease.

13. A method according to any one of claims 1-12, wherein the additional compound is selected from the group consisting of compounds which directly or indirectly interfere with T-cell growth.

14. A method according to claim 13, wherein the compound enhances or inhibits the growth of a certain subgroup of T-cells, such as inflammatory, regulatory or cytotoxic T-cells.

15. A method according to claim 13 or claim 14, wherein the compound is selected from the group consisting of cyclosporin, GM-CSF, Prednisone, Tacrolimus, FK506, IL-10, anti-IL-10, TNF $\alpha$  antibody, IL-12, anti-IL-12, IL-7, anti-IL-7, IL-9, anti-IL-9, IL-16, caspase inhibitors, and functionally similar compounds.

16. A method according to any one of claims 1-15 further comprising a selection procedure.

17. A method according to any one of the claims 1-16, wherein disease associated inflammatory T-cells are expanded and selected.

18. A method according to claim 17, wherein the inflammatory T-cells are cells having a CD4+ phenotype and a type 1 cytokine profile.

19. A method according to claim 18, wherein the inflammatory T-cells are cells contributing in a type 1 inflammatory infiltrate producing IFN $\gamma$  and TNF $\alpha$ .

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

20. A method according to claim 18 or claim 19, wherein the one or more additional compounds is selected from cyclosporine, Prednisone, Tacrolimus, FK506, GM-CSF, IL-12, IL-16, anti-IL-10, anti-TNF $\alpha$ , and functionally similar compounds.

21. A method according to claim 17, wherein the inflammatory T-cells are cells having a CD4+ phenotype and a type 2 cytokine profile.

22. A method according to claim 21, wherein the inflammatory T-cells are cells contributing in a type 2 inflammatory infiltrate producing IL-4 or IL-5.

23. A method according to claim 21 or claim 22, wherein the one or more additional compound is selected from cyclosporine, Prednisone, Tacrolimus, FK506, GM-CSF, IL-16, anti-IL-12, and functionally similar compounds.

24. A method according to any one of claims 17-23, wherein the disease is mediated or partially mediated by type 1 or type 2 inflammatory T-cells, such as chronic inflammatory bowel diseases, for example Crohn's disease and ulcerative colitis, sclerosis, type I diabetes, rheumatoid arthritis, psoriasis, atopic dermatitis, and asthma.

25. A method according to any one of the claims 1-16, wherein disease associated regulatory T-cells are expanded and selected.

26. A method according to claim 25, wherein the regulatory T-cells are cells having a CD4+ phenotype and a type 1 cytokine profile regulating a type 2 inflammatory disease.

5

27. A method according to claim 26, wherein the regulatory T-cells are cells producing INF $\gamma$ .

10

28. A method according to claim 26 or claim 27, wherein the one or more additional compounds is selected from IL-12 and functionally similar compounds.

15

29. A method according to any one of claims 25-28, wherein the disease is mediated or partly mediated by type 2 inflammatory T-cells such as asthma or atopic dermatitis.

20

30. A method according to claim 25, wherein the regulatory T-cells are cells having a CD4 $^{+}$  phenotype and a type 2 cytokine profile regulating a type 1 inflammatory disease.

25

31. A method according to claim 30, wherein the regulatory T-cells are cells producing IL-10 and/or IL-4.

30

20

32. A method according to claim 30 or 31, wherein the one or more additional compounds is selected from anti-IL-12, IL-10, GM-CSF, IL-16, and functionally similar compounds.

35

33. A method according to any of claims 30-32, wherein the disease is mediated or partially mediated by type 1 inflammatory T-cells, such as chronic inflammatory bowel diseases, for example Crohn's disease and ulcerative colitis, sclerosis, type I diabetes, rheumatoid arthritis, and psoriasis.

40

45

34. A method according to any one of the claims 1-16, wherein disease associated cytotoxic T-cells are expanded and selected.

50

35

55

5 35. A method according to claim 34, wherein the cytotoxic  
T-cells are cells having a CD8+ phenotype.

10 36. A method according to claim 34 or claim 35, wherein  
5 the cytotoxic T-cells are tumour infiltrating lymphocytes  
(TIL) or cells having similar properties.

15 37. A method according to any of claims 34-35, wherein  
the one or more additional compounds is selected from GM-  
20 CSF, caspase inhibitors such as Z-VAD,  $\alpha$ -CD95, IL-10, IL-  
12, IL-16, and functionally similar compounds.

25 38. A method according to any of claims 34-38, wherein  
the disease is of neoplastic origin.

30 39. A method according to any one of claims 34-38 wherein  
the disease is malignant melanoma, renal carcinoma,  
breast cancer, lung cancer, cancer of the uterus,  
prostatic cancer, hepatic carcinoma, or cutaneous  
20 lymphoma.

35 40. A continuous T-cell line obtainable by a method  
according to any of claims 1-39.

40 41. A T-cell line according to claim 40, wherein the T-  
cells are inflammatory T-cells.

45 42. A T-cell line according to claim 40, wherein the T-  
cells are regulatory T-cells.

50 43. A T-cell line according to claim 40, wherein the T-  
cells are cytotoxic T-cells.

55 44. A pharmaceutical composition comprising activated  
35 disease associated T-cells prepared according to the  
method of any one of claims 1-39 or a T-cell line

5  
10 according to any one of claims 40-43, optionally comprising one or more pharmaceutically acceptable drugs and/or excipients.

15 5 45. A composition according to claim 44 wherein the T-cells are inflammatory T-cells, regulatory T-cells or cytotoxic T-cells.

20 10 46. A composition according to claim 44 or claim 45, wherein T-cells prepared according to any of the claim 1-39 or a cell line according to claims 40-43 are re-activated in the presence of one or more antigens.

25 15 47. A composition according to claim 46, wherein the antigen or antigens is/are disease associated antigen(s), alloantigen(s) or super-antigen(s).

30 20 48. A composition according to claim 47, wherein the super-antigens are selected from SEA, SEB, SEC, SED, SEE, TSST, Streptococcus pyogenes enterotoxin A, B and C, and Mycoplasma arthritidis antigen.

35 49. A composition according to any one of claims 44-48, wherein the T-cells have been attenuated.

40 25 50. A vaccine comprising activated disease associated inflammatory T-cells prepared according to the method of any one of claims 1-24, a T-cell line according to claim 41 or a composition according to any one of claims 44-48.

45 30 51. A vaccine according to claim 50, wherein T-cells are re-activated in the presence of one or more antigens.

50 35 52. A vaccine according to claim 51, wherein the antigen or antigens is disease associated antigen(s), alloantigen(s) or super-antigen(s).

53. A vaccine according to claim 52, wherein the  
superantigens are selected from SEA, SEB, SEC, SED, SEE,  
TSST, Streptococcus pyogenes enterotoxin A, B and C, and  
Mycoplasma arthritidis antigen.

54. A vaccine according to any one of claims 50-53,  
wherein the T-cells have been attenuated.

55. A pharmaceutical composition for use in an adjuvant  
treatment of a disease comprising disease associated  
regulatory or cytotoxic T-cells prepared according to the  
method of any of claims 1-24, a T-cell line according to  
claim 42 or claim 43 or a composition according to any of  
claims 44-49.

56. A pharmaceutical composition according to claim 55  
for use in treating diseases of inflammatory, auto-  
immune, allergic, neoplastic or transplantation-related  
origin, or combinations thereof.

57. A pharmaceutical composition according to claim 55 or  
claim 56, wherein the T-cells are re-activated in the  
presence of one or more antigens.

58. A pharmaceutical composition according to claim 57,  
wherein the antigen or antigens is disease associated  
antigen(s), alloantigen(s) or super-antigen(s).

59. A pharmaceutical composition according to claim 58,  
wherein the superantigens are selected from SEA, SEB,  
SEC, SED, SEE, TSST, Streptococcus pyogenes enterotoxin  
A, B and C, and Mycoplasma arthritidis antigen.

5  
10 60. A pharmaceutical composition according to any one of claims 55-59, wherein the T-cells are formulated for administration in activated form.

15 61. Use of a T-cell line according to any of claims 40-43, or T-cells prepared according to any of the claims 1-39 in the preparation of a medicament for the treatment of a T-cell associated disease.

20 62. Use according to claim 61, wherein the disease is a disease of inflammatory, auto-immune, allergic, neoplastic or transplantation-related origin, or combinations thereof.

25 63. Use according to claim 62, wherein the disease is an inflammatory bowel disease, such as Crohn's disease and Ulcerative colitis, sclerosis, type I diabetes, rheumatoid arthritis, psoriasis, atopic dermatitis, asthma, malignant melanoma, renal carcinoma, breast  
30 cancer, lung cancer, cancer of the uterus, prostatic cancer, cutaneous lymphoma, asthma, rejection-related disease, or Graft-versus-host-related disease.

35 64. A method for the diagnosis of a disease in a mammal, comprising

40 (a) obtaining a tissue sample from a mammal including a human being, the sample comprising activated T-cells, antigen presenting cells and antigen(s),

45 30 (b) culturing said tissue sample or said activated T-cells in the presence of two or more T-cell growth factors and optionally one or more additional  
50 compound,

35



5

10

(c) observing the presence and/or function of disease associated T-cells, and relating the presence and/or functional characteristics of these T-cells to a disease.

5

15

65. A method according to claim 64, wherein the disease is related to the disease associated T-cells by determining the kind of activated T-cells and/or their state of activation.

10

20

66. A method according to claim 64 or 65, wherein the cytokine profile of the T-cells is determined.

25

15

67. A method for the treatment, alleviation or prevention of a disease associated with an activation of T-cells in a subject comprising administering a T-cell line according to any of claims 40-43, T-cells as produced according to any of claims 1-39, a composition according to any of claims 44-49 or 5-58, or a vaccine according to claims 50-54 to said subject.

30

20

35

68. A method according to claim 67, wherein the T-cells are expanded from a tissue sample collected from the patient to be treated.

25

40

69. A method according to claim 67, wherein the T-cells are expanded from a tissue sample collected from a patient different to the patient to be treated.

45

30

70. A method according to claim 69, further comprising determining the HLA restriction in the T-cells and in the patient to be treated.

50

35

71. A method of testing the effect of a medicament against a T-cell associated disease comprising

55

5

- (a) providing a T-cell line according to any of claims 39-42,
- 10 (b) applying the medicament to be tested to the T-cell line, and
- 5 (c) observing the effect of the medicament on the T-cell line.

15

72. A method according to claim 71, wherein the cytokine profile of the T-cell line with and without the addition of the medicament is compared.

20

73. A method according to claim 71, wherein the phenotype, proliferation and/or apoptosis of the T-cell line with and without the addition of the medicament is compared.

25

74. A method according to claim 71, wherein the intracellular amount of NFkB and/or JAK/STAT pathway is/are monitored.

30

20

75. A method according to any of claims 71-73, wherein the medicament to be tested is selected from compound libraries such as small molecule libraries or peptide libraries or antibodies against T-cell components.

35

25

76. A method according to any of claims 71-75, wherein the medicament is selected from peptide fragments from T-cell receptors.

40

77. A model system for testing the effect of a medicament against a T-cell associated disease comprising at least one T-cell line according to any one of claims 40-43.

45

78. A method for the treatment, alleviation or prevention of a disease associated with an activation of T-cells in a subject comprising administering a medicament as

50

55

identified according to the method of claims 71-76 as being effective in said treatment.

79. A method according to claim 78 wherein the disease is a disease of inflammatory, auto-immune, allergic, neoplastic or transplantation-related origin, or combinations thereof.

80. A method according to claim 78 or claim 79, wherein the disease is an inflammatory bowel disease such as Crohn's colitis or ulcerative colitis, sclerosis, type I diabetes, rheumatoid arthritis, psoriasis, atopic dermatitis, asthma, malignant melanoma, renal carcinoma, lung cancer, cancer of the uterus, prostate cancer, hepatic carcinoma, breast cancer, cutaneous lymphoma, rejection-related disease or Graft-versus-host-related disease.

81. A method of detecting T-cell growth factors for use in the method according to any of claims 1-39, wherein candidate factors are tested in a method according to claim 5 in place of IL-2 or IL-4 or a functionally similar compound or in addition to the combination of IL-2 and IL-4 or said functionally similar compound(s), and the effect compared to the effect obtained by using a combination of IL-2 and IL-4.

82. A method of monitoring the response to a treatment of a disease of inflammatory, auto-immune, allergic, neoplastic or transplantation-related origin, or combinations thereof, comprising comparing the phenotype, proliferation, apoptosis, cytokine profile, intracellular amount of NFkB and/or JAK/STAT pathway of activated T-cells in tissue sample taken from the patient to be treated before the start of the treatment and during the treatment and/or after the treatment has ended.

5

10

83. A method according to claim 82, wherein patients which do not respond to a certain treatment are identified.

5

15

84. A method of identifying disease associated antigens, comprising screening peptide libraries or antigen samples for their re-activation properties in a T-cell line according to any of claims 40-43.

10

20

25

30

35

40

45

50

55

# T-cell vaccination - principles




FIG. 1

# T-cell culture

**100-500 x 10<sup>6</sup>**

## T - cells


RPMI 1640  
IL-2+IL 4

2/22

Clonality - 2-3 clones

**Inactivation by  $\gamma$  - irradiation**

FIG. 2



# T - cell vaccination - procedure





FIG. 3

4/22

FIG. 4





5/22





FIG. 5


6/22

FIG. 6




7/22

FIG. 7




8/22

FIG. 8




9/22

FIG. 9




10/22

FIG. 10




11/22

FIG. 11



12/22


FIG. 12





13/22

FIG. 13




# Best Available Copy

WO 00/00587

PCT/DK99/00363

14/22

FIG. 14




# Best Available Copy

WO 00/00587

PCT/DK99/00363

15/22


FIG. 15



# Best Available Copy


WO 00/00587

PCT/DK99/00363




17/22

FIG. 17




18/22


FIG. 18




Quad	Events	% Gated	% Total	X Mean	Y Mean
UL	51	0.51	0.51	63.72	327.75
UR	3483	34.83	34.83	715.95	429.73
LL	5554	55.54	55.54	16.04	1.58
LR	912	9.12	9.12	554.74	21.80




Quad	Events	% Gated	% Total	X Mean	Y Mean
UL	5	0.05	0.05	33.83	272.70
UR	664	6.64	6.64	1544.52	544.57
LL	8303	83.03	83.03	15.61	1.24
LR	1028	10.28	10.28	968.75	25.23




Quad	Events	% Gated	% Total	X Mean	Y Mean
UL	559	5.59	5.59	60.59	420.74
UR	1915	19.15	19.15	399.96	512.10
LL	5410	54.10	54.10	33.57	4.28
LR	2116	21.16	21.16	346.08	16.63



Quad	Events	% Gated	% Total	X Mean	Y Mean
UL	2	0.02	0.02	36.30	252.48
UR	760	7.60	7.60	1151.09	446.68
LL	7829	78.29	78.29	17.73	1.39
LR	1409	14.09	14.09	707.89	20.83



Quad	Events	% Gated	% Total	X Mean	Y Mean
UL	14	0.14	0.14	67.28	226.32
UR	388	3.88	3.88	1058.27	402.01
LL	8744	87.44	87.44	11.86	1.31
LR	854	8.54	8.54	783.76	21.95



Quad	Events	% Gated	% Total	X Mean	Y Mean
UL	454	4.54	4.54	85.27	513.96
UR	2815	28.15	28.15	483.48	647.67
LL	4615	46.15	46.15	41.28	4.54
LR	2116	21.16	21.16	407.90	20.88


19/22

C 8.3

Test Number 00000200  
Analysis Time: 24-Feb-99 08:38:22  
Time Printed: 24-Feb-99 08:38:38  
Version 22 Version 1.01

Instrument Settings  
Aperture Diameter: 100um C  
Kd: 56.83  
Metered Volume: 0.5 ml  
Dilution Factor: 1  
Repetitions: 1 runs  
Amplifier, Pre-Amp Gain: 224.00  
Amplifier, Main Gain: 64  
Aperture Current: 0.354 mA

Counting Results  
Count > 5.000 um: 14042  
Count > 15.34 um: 251  
Count Between: 13791



Cursor Statistics  
Size at C1: 6.774 um  
Size at C2: 12.28 um  
Mean: 9.860  
Median: 9.891  
Mode: 9.109  
Standard Deviation: 1.207  
Channelizer Count: 11042

Statistical Results  
Mean: 9.533  
Median: 9.745  
Mode: 4.934  
Std Dev: 2.135

FIG. 19A

20/22

C8.2


Test Number 00000207  
Analysis Time: 25-Feb-99 08:50:11  
Time Printed: 25-Feb-99 08:50:26  
Version Z2 Version 1.01

Instrument Settings

Aperture Diameter: 100um C  
Kd: 56.83  
Metered Volume: 0.5 ml  
Dilution Factor: 1  
Repetitions: 1 runs  
Amplifier, Pre-Amp Gain: 224.00  
Amplifier, Main Gain: 64  
Aperture Current: 0.354 mA

Counting Results

Count > 5.000 um: 18264  
Count > 15.34 um: 157  
Count Between: 18107



Cursor Statistics  
Size at C1: 6.774 um  
Size at C2: 12.28 um  
Mean: 9.823  
Median: 9.843  
Mode: 9.542  
Standard Deviation: 1.160  
Channelizer Count: 14722

Statistical Results  
Mean: 9.394  
Median: 9.644  
Mode: 4.934  
Std Dev: 2.041

FIG. 19B




21/22

C8.3 + inf1.

Test Number 00000208  
 Analysis Time: 25-Feb-99 08:51:30  
 Time Printed: 25-Feb-99 08:51:46  
 Version Z2 Version 1.01

Instrument Settings  
 Aperture Diameter: 100um C  
 Kd: 56.83  
 Metered Volume: 0.5 ml  
 Dilution Factor: 1  
 Repetitions: 1 runs  
 Amplifier, Pre-Amp Gain: 224.00  
 Amplifier, Main Gain: 64  
 Aperture Current: 0.354 mA

Counting Results  
 Count > 5.000 um: 17198  
 Count > 15.34 um: 107  
 Count Between: 17091



Cursor Statistics  
 Size at C1: 6.774 um  
 Size at C2: 12.28 um  
 Mean: 9.944  
 Median: 9.939  
 Mode: 9.644  
 Standard Deviation: 1.160  
 Channelizer Count: 14111

Statistical Results  
 Mean: 9.608  
 Median: 9.843  
 Mode: 4.934  
 Std Dev: 2.008

FIG. 19C



FIG. 20A



FIG. 20B

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/DK 99/00363

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>		
IPC 6	C12N5/06	C12N5/08 A61K39/00 A61K35/14 G01N33/50
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b>		
Minimum documentation searched (classification system followed by classification symbols)		
IPC 6 C12N A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	K. KALTOFT ET AL.: "COMMON CLONAL CHROMOSOME ABERRATIONS IN CYTOKINE-DEPENDENT CONTINUOUS HUMAN T-LYMPHOCYTE CELL LINES." CANCER GENETICS AND CYTOGENETICS, vol. 85, no. 1, November 1995 (1995-11), pages 69-71, XP002087068 NEW YORK, N.Y., US cited in the application the whole document --- -/-	1-12, 40
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
5 October 1999		18/10/1999
Name and mailing address of the ISA European Patent Office, P.B. 5018 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo ni, Fax: (+31-70) 340-3016		Authorized officer  Ryckebosch, A

Form PCT/ISA/210 (second sheet) (July 1992)

# INTERNATIONAL SEARCH REPORT

Int: onal Application No

PCT/DK 99/00363

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	K. KALTOFT ET AL.: "IN VITRO GENETICALLY ABERRANT T-CELL CLONES WITH CONTINUOUS GROWTH ARE ASSOCIATED WITH ATOPIC DERMATITIS." ARCHIVES OF DERMATOLOGICAL RESEARCH, vol. 287, no. 1, December 1994 (1994-12), pages 42-47, XP002087069 CHICAGO, IL, US cited in the application the whole document	1-12,40
A	R.B. EFFROS ET AL.: "REPLICATIVE SENESCENCE OF T CELLS: DOES THE HAYFLICK LIMIT LEAD TO IMMUNE EXHAUSTION?" IMMUNOLOGY TODAY, vol. 18, no. 9, September 1997 (1997-09), pages 450-454, XP002087067 AMSTERDAM, NL the whole document	1-63, 67-70
A	WO 94 02156 A (THE BOARD OF TRUSTEES OF LELAND STANFORD JUNIOR UNIVERSITY) 3 February 1994 (1994-02-03) cited in the application page 8, line 30 -page 10, line 6; claims 1-11; example 7 page 18, line 35 -page 19, line 12 page 21, line 1 -page 24, line 21	1-63, 67-70
A	WO 97 05239 A (CELLTHERAPY, INC.) 13 February 1997 (1997-02-13) cited in the application page 11, line 10 -page 16, line 26; claims	1-63, 67-70
A	WO 88 07077 A (THE CHILDREN'S HOSPITAL, INC.) 22 September 1988 (1988-09-22) cited in the application claims	1-63, 67-70
P,X	K. KALTOFT: "CYTOKINE-DRIVEN IMMORTALIZATION OF IN VITRO ACTIVATED HUMAN T LYMPHOCYTES: CD28 EXPRESSION CORRELATES INVERSELY WITH CELL POPULATION DOUBLINGS." EXPERIMENTAL AND CLINICAL IMMUNOGENETICS, vol. 15, no. 2, July 1998 (1998-07), pages 84-89, XP002117587 BASEL, CH the whole document	1-84

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 99/00363

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 67-70 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 78-80  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 78-80

Claims 78-80, relating to a method for treatment which comprises administering a medicament as identified according to a testing method of claims 71-76, lack any structural feature which could be the basis for a meaningful search and consequently have not been searched as their subject-matters were not sufficiently disclosed and lack adequate support in the description (Art. 5 and 6 PCT).

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/DK 99/00363

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
W0 9402156	A	03-02-1994	AU 4678993 A	14-02-1994
W0 9705239	A	13-02-1997	AU 6649996 A	26-02-1997
			CA 2227327 A	13-02-1997
			EP 0852618 A	15-07-1998
W0 8807077	A	22-09-1988	AU 1363088 A	10-10-1988
			DK 628088 A	11-01-1989
			EP 0348413 A	03-01-1990
			JP 2502424 T	09-08-1990

***This Page Blank (uspto)***



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

*This Page Blank (uspto)*